


Accepted Manuscript



2013 ACC/AHA Guideline on the Assessment of Cardiovascular Risk

David C. Goff, MD, PhD, FACP, FAHA Donald M. Lloyd-Jones, MD, ScM, FACC, FAHA Glen Bennett, MPH Christopher J. O'Donnell, MD, MPH Sean Coady, MS Jennifer Robinson, MD, MPH, FAHA Ralph B. D'Agostino, J. Sanford Schwartz, MD Raymond Gibbons, MD, FACC, FAHA Susan T. Shero, MS, RN Philip Greenland, MD, FACC, FAHA Sidney C. Smith, MD, FACC, FAHA Daniel T. Lackland, DrPH, FAHA Paul Sorlie, PhD Daniel Levy, MD Neil J. Stone, MD, FACC, FAHA Peter W.F. Wilson, MD, FAHA

PII: S0735-1097(13)06031-2

DOI: [10.1016/j.jacc.2013.11.005](https://doi.org/10.1016/j.jacc.2013.11.005)

Reference: JAC 19599

To appear in: *Journal of the American College of Cardiology*

Please cite this article as: Goff Jr DC, Lloyd-Jones DM, Bennett G, O'Donnell CJ, Coady S, Robinson J, D'Agostino Sr RB, Schwartz JS, Gibbons R, Shero ST, Greenland P, Smith Jr SC, Lackland DT, Sorlie P, Levy D, Stone NJ, Wilson PWF, 2013 ACC/AHA Guideline on the Assessment of Cardiovascular Risk, *Journal of the American College of Cardiology* (2013), doi: 10.1016/j.jacc.2013.11.005.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Goff DC Jr, et al.
2013 ACC/AHA Cardiovascular Risk Guideline

2013 ACC/AHA Guideline on the Assessment of Cardiovascular Risk

A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines

Endorsed by the American Association of Cardiovascular and Pulmonary Rehabilitation, American Society for Preventive Cardiology, American Society of Hypertension, Association of Black Cardiologists, National Lipid Association, Preventive Cardiovascular Nurses Association, and WomenHeart: The National Coalition for Women with Heart Disease

EXPERT WORK GROUP MEMBERS

David C. Goff, Jr, MD, PhD, FACP, FAHA, *Co-Chair*
Donald M. Lloyd-Jones, MD, ScM, FACC, FAHA, *Co-Chair*
Glen Bennett, MPH*
Sean Coady, MS*
Ralph B. D'Agostino, Sr, PhD, FAHA
Raymond Gibbons, MD, FACC, FAHA
Philip Greenland, MD, FACC, FAHA
Daniel T. Lackland, DrPH, FAHA
Daniel Levy, MD*
Christopher J. O'Donnell, MD, MPH*
Jennifer Robinson, MD, MPH, FAHA
J. Sanford Schwartz, MD
Susan T. Shero, MS, RN*
Sidney C. Smith, Jr, MD, FACC, FAHA
Paul Sorlie, PhD*
Neil J. Stone, MD, FACC, FAHA
Peter W.F. Wilson, MD, FAHA

Methodology Members

Harmon S. Jordan, ScD
Lev Nevo, MD
Janusz Wnek, PhD

ACCF/AHA TASK FORCE MEMBERS

Jeffrey L. Anderson, MD, FACC, FAHA, *Chair*
Jonathan L. Halperin, MD, FACC, FAHA, *Chair-Elect*
Nancy M. Albert, PhD, CCNS, CCRN, FAHA
Biykem Bozkurt, MD, PhD, FACC, FAHA
Ralph G. Brindis, MD, MPH, MACC
Lesley H. Curtis, PhD, FAHA
David DeMets, PhD
Robert A. Guyton, MD, FACC
Judith S. Hochman, MD, FACC, FAHA
Richard J. Kovacs, MD, FACC, FAHA
E. Magnus Ohman, MD, FACC
Susan J. Pressler, PhD, RN, FAAN, FAHA
Frank W. Sellke, MD, FACC, FAHA
Win-Kuang Shen, MD, FACC, FAHA

Subcommittee on Prevention Guidelines

Sidney C. Smith, Jr, MD, FACC, FAHA, *Chair*
Gordon F. Tomaselli, MD, FACC, FAHA, *Co-Chair*

*Ex-Officio Members.

This document was approved by the American College of Cardiology Board of Trustees and the American Heart Association Science Advisory and Coordinating Committee in November 2013.

The American College of Cardiology requests that this document be cited as follows: Goff DC Jr, Lloyd-Jones DM, Bennett G, Coady S, D'Agostino RB Sr, Gibbons R, Greenland P, Lackland DT, Levy D, O'Donnell CJ, Robinson J, Schwartz JS, Smith SC Jr, Sorlie P, Shero ST, Stone NJ, Wilson PW. 2013 ACC/AHA guideline on the assessment of cardiovascular risk: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol* 2013; ●●:●●●●-●●●●.

This article is copublished in *Circulation*.

Goff DC Jr, et al.
2013 ACC/AHA Cardiovascular Risk Guideline

Copies: This document is available on the World Wide Web sites of the American College of Cardiology (www.cardiosource.org), and the American Heart Association (my.americanheart.org). A copy of the document is available at <http://my.americanheart.org/statements> by selecting either the "By Topic" link or the "By Publication Date" link. For copies of this document, please contact the Elsevier Inc. Reprint Department, fax (212) 633-3820, e-mail reprints@elsevier.com.

Permissions: Multiple copies, modification, alteration, enhancement, and/or distribution of this document are not permitted without the express permission of the American College of Cardiology Foundation. Please contact healthpermissions@elsevier.com.

©2013 The Expert Work Group Members. The *Journal of the American College of Cardiology* is published on behalf of the American College of Cardiology Foundation by Elsevier Inc.; *Circulation* is published on behalf of the American Heart Association, Inc., by Wolters Kluwer. This is an open access article under the terms of the Creative Commons Attribution Non-Commercial-NoDerivs License, which permits use, distribution, and reproduction in any medium, provided that the Contribution is properly cited, the use is non-commercial, and no modifications or adaptations are made.

Table of Contents

Preamble and Transition to ACC/AHA Guidelines to Reduce Cardiovascular Risk.....	4
1. Introduction	9
1.1. Organization of the Work Group	9
1.2. Document Review and Approval.....	9
1.3. Charge to the Work Group.....	9
1.4. Methodology and Evidence Review	9
2. Risk Assessment: Recommendations	10
3. Approach to Risk Assessment	11
4. Development of New Pooled Cohort ASCVD Risk Equations	13
4.1. Recommendations for Assessment of 10-Year Risk for a First Hard ASCVD Event.....	14
5. Implications for Risk Assessment.....	15
6. CQs and Systematic Evidence Review	18
6.1. Critical Question 1	18
6.1.1. Summary of Systematic Reviews and Meta-Analyses for CQ1	19
6.1.2. Recommendations for CQ1: use of Newer Risk Markers After Quantitative Risk Assessment	20
6.2. Critical Question 2	21
6.2.1. Summary of Evidence for CQ2	21
6.2.2. Recommendations for CQ2: Long-Term Risk Assessment.....	22
7. Implementation Considerations for Risk Assessment	23
8. Evidence Gaps and Future Research Needs.....	24
9. Conclusions	24
Appendix 1. Evidence Statements for CQ1	26
Appendix 2. Evidence Statements for CQ2.....	32
Appendix 3. Characteristics of Previously Published Risk Scores and Current Pooled Cohort Equations	34
Appendix 4. Development and Steps for Implementation of the ASCVD Pooled Cohort Risk Equations	36
Appendix 5. Author Relationships With Industry and Other Entities (Relevant).....	41
Appendix 6. ACC/AHA Expert Reviewer Relationships With Industry and Other Entities	44
Appendix 7. Abbreviations.....	45
References	46

Preamble and Transition to ACC/AHA Guidelines to Reduce Cardiovascular Risk

The goals of the American College of Cardiology (ACC) and the American Heart Association (AHA) are to prevent cardiovascular (CV) diseases, improve the management of people who have these diseases through professional education and research, and develop guidelines, standards and policies that promote optimal patient care and CV health. Toward these objectives, the ACC and AHA have collaborated with the National Heart, Lung, and Blood Institute (NHLBI) and stakeholder and professional organizations to develop clinical practice guidelines for assessment of CV risk, lifestyle modifications to reduce CV risk, and management of blood cholesterol, overweight and obesity in adults.

In 2008, the NHLBI initiated these guidelines by sponsoring rigorous systematic evidence reviews for each topic by expert panels convened to develop critical questions (CQs), interpret the evidence and craft recommendations. In response to the 2011 report of the Institute of Medicine on the development of trustworthy clinical guidelines (1), the NHLBI Advisory Council (NHLBAC) recommended that the NHLBI focus specifically on reviewing the highest quality evidence and partner with other organizations to develop recommendations (2,3). Accordingly, in June 2013 the NHLBI initiated collaboration with the ACC and AHA to work with other organizations to complete and publish the 4 guidelines noted above and make them available to the widest possible constituency. Recognizing that the expert panels did not consider evidence beyond 2011 (except as specified in the methodology), the ACC, AHA, and collaborating societies plan to begin updating these guidelines starting in 2014.

The joint ACC/AHA Task Force on Practice Guidelines (Task Force) appointed a subcommittee to shepherd this transition, communicate the rationale and expectations to the writing panels and partnering organizations and expeditiously publish the documents. The ACC/AHA and partner organizations recruited a limited number of expert reviewers for fiduciary examination of content, recognizing that each document had undergone extensive peer review by representatives of the NHLBAC, key Federal agencies and scientific experts. Each writing panel responded to comments from these reviewers. Clarifications were incorporated where appropriate, but there were no substantive changes as the bulk of the content was undisputed.

Although the Task Force led the final development of these prevention guidelines, they differ from other ACC/AHA guidelines. First, as opposed to an extensive compendium of clinical information, these documents are significantly more limited in scope and focus on selected CQs in each topic, based on the highest quality evidence available. Recommendations were derived from randomized trials, meta-analyses, and observational studies evaluated for quality, and were not formulated when sufficient evidence was not available. Second, the text accompanying each recommendation is succinct, summarizing the evidence for each question. The Full Panel Reports include more detailed information

about the evidence statements that serves as the basis for recommendations. Third, the format of the recommendations differs from other ACC/AHA guidelines. Each recommendation has been mapped from the NHLBI grading format to the ACC/AHA Class of Recommendation/Level of Evidence (COR/LOE) construct (Table 1) and is expressed in both formats. Because of the inherent differences in grading systems and the clinical questions driving the recommendations, alignment between the NHLBI and ACC/AHA formats is in some cases imperfect. Explanations of these variations are noted in the recommendation tables, where applicable.

Table 1. Applying Classification of Recommendation and Level of Evidence

		SIZE OF TREATMENT EFFECT												
		CLASS I <i>Benefit >>> Risk</i> Procedure/Treatment SHOULD be performed/ administered	CLASS IIa <i>Benefit >> Risk</i> Additional studies with <i>focused objectives</i> needed IT IS REASONABLE to per- form procedure/administer treatment	CLASS IIb <i>Benefit ≥ Risk</i> Additional studies with <i>broad objectives</i> needed; additional registry data would be <i>helpful</i> Procedure/Treatment MAY BE CONSIDERED	CLASS III <i>No Benefit</i> or CLASS III <i>Harm</i> <table><tr><th colspan="2">Procedure/ Test</th><th>Treatment</th></tr><tr><td>COR III: No benefit</td><td>Not Helpful</td><td>No Proven Benefit</td></tr><tr><td>COR III: Harm</td><td>Excess Cost w/o Benefit or Harmful</td><td>Harmful to Patients</td></tr></table>	Procedure/ Test		Treatment	COR III: No benefit	Not Helpful	No Proven Benefit	COR III: Harm	Excess Cost w/o Benefit or Harmful	Harmful to Patients
Procedure/ Test		Treatment												
COR III: No benefit	Not Helpful	No Proven Benefit												
COR III: Harm	Excess Cost w/o Benefit or Harmful	Harmful to Patients												
ESTIMATE OF CERTAINTY (PRECISION) OF TREATMENT EFFECT	LEVEL A Multiple populations evaluated* Data derived from multiple randomized clinical trials or meta-analyses	<ul style="list-style-type: none">■ Recommendation that procedure or treatment is useful/effective■ Sufficient evidence from multiple randomized trials or meta-analyses	<ul style="list-style-type: none">■ Recommendation in favor of treatment or procedure being useful/effective■ Some conflicting evidence from multiple randomized trials or meta-analyses	<ul style="list-style-type: none">■ Recommendation's usefulness/efficacy less well established■ Greater conflicting evidence from multiple randomized trials or meta-analyses	<ul style="list-style-type: none">■ Recommendation that procedure or treatment is not useful/effective and may be harmful■ Sufficient evidence from multiple randomized trials or meta-analyses									
	LEVEL B Limited populations evaluated* Data derived from a single randomized trial or nonrandomized studies	<ul style="list-style-type: none">■ Recommendation that procedure or treatment is useful/effective■ Evidence from single randomized trial or nonrandomized studies	<ul style="list-style-type: none">■ Recommendation in favor of treatment or procedure being useful/effective■ Some conflicting evidence from single randomized trial or nonrandomized studies	<ul style="list-style-type: none">■ Recommendation's usefulness/efficacy less well established■ Greater conflicting evidence from single randomized trial or nonrandomized studies	<ul style="list-style-type: none">■ Recommendation that procedure or treatment is not useful/effective and may be harmful■ Evidence from single randomized trial or nonrandomized studies									
	LEVEL C Very limited populations evaluated* Only consensus opinion of experts, case studies, or standard of care	<ul style="list-style-type: none">■ Recommendation that procedure or treatment is useful/effective■ Only expert opinion, case studies, or standard of care	<ul style="list-style-type: none">■ Recommendation in favor of treatment or procedure being useful/effective■ Only diverging expert opinion, case studies, or standard of care	<ul style="list-style-type: none">■ Recommendation's usefulness/efficacy less well established■ Only diverging expert opinion, case studies, or standard of care	<ul style="list-style-type: none">■ Recommendation that procedure or treatment is not useful/effective and may be harmful■ Only expert opinion, case studies, or standard of care									
	Suggested phrases for writing recommendations		should is recommended is indicated is useful/effective/beneficial	is reasonable can be useful/effective/beneficial is probably recommended or indicated	may/might be considered may/might be reasonable usefulness/effectiveness is unknown/unclear/uncertain or not well established	COR III: No Benefit is not recommended is not indicated should not be performed/ administered/ other is not useful/ beneficial/ effective COR III: Harm potentially harmful causes harm associated with excess morbid ity/mortality should not be performed/ administered/ other								
Comparative effectiveness phrases [†]		treatment/strategy A is recommended/indicated in preference to treatment B treatment A should be chosen over treatment B	treatment/strategy A is probably recommended/indicated in preference to treatment B it is reasonable to choose treatment A over treatment B											

A recommendation with Level of Evidence B or C does not imply that the recommendation is weak. Many important clinical questions addressed in the guidelines do not lend themselves to clinical trials. Even when

Goff DC Jr, et al.
2013 ACC/AHA Cardiovascular Risk Guideline

randomized trials are unavailable, there may be a very clear clinical consensus that a particular test or therapy is useful or effective.

*Data available from clinical trials or registries about the usefulness/efficacy in different subpopulations, such as sex, age, history of diabetes, history of prior myocardial infarction, history of heart failure, and prior aspirin use.

†For comparative effectiveness recommendations (Class I and IIa; Level of Evidence A and B only), studies that support the use of comparator verbs should involve direct comparisons of the treatments or strategies being evaluated.

In consultation with NHLBI, the policies adopted by the writing panels to manage relationships of authors with industry and other entities (RWI) are outlined in the methods section of each panel report. These policies were in effect when this effort began in 2008 and throughout the writing process and voting on recommendations, until the process was transferred to ACC/AHA in 2013. In the interest of transparency, the ACC/AHA requested that panel authors resubmit RWI disclosures as of July 2013. Relationships relevant to this guideline are disclosed in Appendix 5. None of the ACC/AHA expert reviewers had relevant RWI (Appendix 6).

Systematic evidence reports and accompanying summary tables were developed by the expert panels and NHLBI. The guideline was reviewed by the ACC/AHA Task Force and approved by the ACC Board of Trustees, the AHA Science Advisory and Coordinating Committee, and the governing bodies of partnering organizations. In addition, ACC/AHA sought endorsement by other stakeholders, including professional organizations. It is the hope of the writing panels, stakeholders, professional organizations, NHLBI, and the Task Force that the guidelines will garner the widest possible readership for the benefit of patients, providers and the public health.

Guidelines attempt to define practices that meet the needs of patients in most circumstances and are not a replacement for clinical judgment. The ultimate decision about care of a particular patient must be made by the healthcare provider and patient in light of the circumstances presented by that patient. As a result, situations might arise in which deviations from these guidelines may be appropriate. These considerations notwithstanding, in caring for most patients, clinicians can employ the recommendations confidently to reduce the risks of atherosclerotic cardiovascular disease (ASCVD) events.

See Tables 2 and 3 for an explanation of the NHLBI recommendation grading methodology.

Table 2. NHLBI Grading the Strength of Recommendations

Grade	Strength of Recommendation*
A	Strong recommendation There is high certainty based on evidence that the net benefit† is substantial.
B	Moderate recommendation There is moderate certainty based on evidence that the net benefit is moderate to substantial, or there is high certainty that the net benefit is moderate.

C	Weak recommendation There is at least moderate certainty based on evidence that there is a small net benefit.
D	Recommendation against There is at least moderate certainty based on evidence that it has no net benefit or that risks/harms outweigh benefits.
E	Expert opinion (“There is insufficient evidence or evidence is unclear or conflicting, but this is what the Work Group recommends.”) Net benefit is unclear. Balance of benefits and harms cannot be determined because of no evidence, insufficient evidence, unclear evidence, or conflicting evidence, but the Work Group thought it was important to provide clinical guidance and make a recommendation. Further research is recommended in this area.
N	No recommendation for or against (“There is insufficient evidence or evidence is unclear or conflicting.”) Net benefit is unclear. Balance of benefits and harms cannot be determined because of no evidence, insufficient evidence, unclear evidence, or conflicting evidence, and the Work Group thought no recommendation should be made. Further research is recommended in this area.

*In most cases, the strength of the recommendation should be closely aligned with the quality of the evidence; however, under some circumstances, there may be valid reasons for making recommendations that are not closely aligned with the quality of the evidence (e.g., strong recommendation when the evidence quality is moderate, like smoking cessation to reduce CVD risk or ordering an ECG as part of the initial diagnostic work-up for a patient presenting with possible MI). Those situations should be limited and the rationale explained clearly by the Work Group.

†Net benefit is defined as benefits minus risks/harms of the service/intervention.

CVD indicates cardiovascular risk; ECG, electrocardiography; MI, myocardial infarction; and NHLBI, National Heart, Lung, and Blood Institute.

Table 3. Quality Rating the Strength of Evidence

Type of Evidence	Quality Rating*
<ul style="list-style-type: none"> Well-designed, well-executed† RCTs that adequately represent populations to which the results are applied and directly assess effects on health outcomes. MAs of such studies. <p>Highly certain about the estimate of effect. Further research is unlikely to change our confidence in the estimate of effect.</p>	High
<ul style="list-style-type: none"> RCTs with minor limitations‡ affecting confidence in, or applicability of, the results. Well-designed, well-executed nonrandomized controlled studies§ and well-designed, well-executed observational studies . MAs of such studies. <p>Moderately certain about the estimate of effect. Further research may have an impact on our confidence in the estimate of effect and may change the estimate.</p>	Moderate

Goff DC Jr, et al.
2013 ACC/AHA Cardiovascular Risk Guideline

<ul style="list-style-type: none"> • RCTs with major limitations. • Nonrandomized controlled studies and observational studies with major limitations affecting confidence in, or applicability of, the results. • Uncontrolled clinical observations without an appropriate comparison group (e.g., case series, case reports). • Physiological studies in humans. • MAs of such studies. <p>Low certainty about the estimate of effect. Further research is likely to have an impact on our confidence in the estimate of effect and is likely to change the estimate.</p>	Low
---	-----

*In some cases, other evidence, such as large all-or-none case series (e.g., jumping from airplanes or tall structures), can represent high or moderate quality evidence. In such cases, the rationale for the evidence rating exception should be explained by the Work Group and clearly justified.

†Well-designed, well-executed refers to studies that directly address the question, use adequate randomization, blinding, allocation concealment, are adequately powered, use ITT analyses, and have high follow-up rates.

‡Limitations include concerns with the design and execution of a study that result in decreased confidence in the true estimate of the effect. Examples of such limitations include, but are not limited to: inadequate randomization, lack of blinding of study participants or outcome assessors, inadequate power, outcomes of interest are not prespecified or the primary outcomes, low follow-up rates, or findings based on subgroup analyses. Whether the limitations are considered minor or major is based on the number and severity of flaws in design or execution. Rules for determining whether the limitations are considered minor or major and how they will affect rating of the individual studies will be developed collaboratively with the methodology team.

§Nonrandomized controlled studies refer to intervention studies where assignment to intervention and comparison groups is not random (e.g., quasi-experimental study design)

|| Observational studies include prospective and retrospective cohort, case-control, and cross sectional studies.

ITT indicates intention-to-treat; MA, meta-analysis; and RCT, randomized controlled trial.

1. Introduction

1.1. Organization of the Work Group

The Risk Assessment Work Group (Work Group) was composed of 11 members and 5 ex-officio members, including internists, cardiologists, endocrinologists, and experts in CV epidemiology, biostatistics, healthcare management and economics, and guideline development.

1.2. Document Review and Approval

A formal peer review process, which included 12 expert reviewers and representatives of Federal agencies, was initially completed under the auspices of the NHLBI. This document was also reviewed by 3 expert reviewers nominated by the ACC and the AHA when the management of the guideline transitioned to the ACC/AHA. The ACC and AHA Reviewers' RWI information is published in this document (6).

This document was approved for publication by the governing bodies of the ACC and AHA and endorsed by the American Association of Cardiovascular and Pulmonary Rehabilitation, American Society for Preventive Cardiology, American Society of Hypertension, Association of Black Cardiologists, National Lipid Association, Preventive Cardiovascular Nurses Association, and WomenHeart: The National Coalition for Women with Heart Disease.

1.3. Charge to the Work Group

The Work Group was 1 of 3 work groups appointed by the NHLBI to develop its own recommendations and provide cross-cutting input to 3 Expert Panels for updating guidelines on blood cholesterol, blood pressure (BP), and overweight/obesity.

The Work Group was asked to examine the scientific evidence on risk assessment for initial ASCVD events, and to develop an approach for risk assessment that could be used in practice and used or adapted by the risk factor panels (cholesterol, hypertension, and obesity) in their guidelines and algorithms. Specifically, the Work Group was charged with 2 tasks:

1. To develop or recommend an approach to quantitative risk assessment that could be used to guide care; and
2. To pose and address a small number of questions judged to be critical to refining and adopting risk assessment in clinical practice using systematic review methodology.

1.4. Methodology and Evidence Review

This guideline is based on the Full Work Group Report which is provided as a supplement to the guideline (http://jaccjacc.cardiosource.com/acc_documents/2013_FPR_S5_Risk_Assesment.pdf). The Full Work Group Report contains background and additional material related to content, methodology, evidence synthesis, rationale, and references and is supported by the NHLBI Systematic Evidence Review

which can be found at (http://www.nhlbi.nih.gov/guidelines/cvd_adult/risk_assessment/). These documents also describe the process for the development of novel, comprehensive multivariable risk equations for the prediction of 10-year risk for development of ASCVD in nonHispanic African-American and nonHispanic White men and women from 40 to 79 years of age. These equations were developed from several long-standing population-based cohort studies funded by the NHLBI. Ten-year risk was defined as the risk of developing a first ASCVD event, defined as nonfatal myocardial infarction or coronary heart disease (CHD) death, or fatal or nonfatal stroke, over a 10-year period among people free from ASCVD at the beginning of the period.

In addition, through evaluation of evidence developed through systematic reviews of the literature, the Work Group addressed the following 2 CQs:

CQ1: “What is the evidence regarding reclassification or contribution to risk assessment when high-sensitivity C-reactive protein (hs-CRP), apolipoprotein B (ApoB), glomerular filtration rate (GFR), microalbuminuria, family history, cardiorespiratory fitness, ankle-brachial index (ABI), carotid intima-media thickness (CIMT), or coronary artery calcium (CAC) score are considered in addition to the variables that are in the traditional risk scores?”

CQ2: “Are models constructed to assess the long-term (≥ 15 years or lifetime) risk for a first cardiovascular disease (CVD) event in adults effective in assessing variation in long-term risk among adults at low and/or intermediate short-term risk, whether analyzed separately or combined?”

The evidence and recommendations in the guideline focus on the large proportion of the adult population without clinical signs or symptoms of ASCVD, who merit evaluation for the primary prevention of ASCVD. They do not apply to those with clinically-manifest ASCVD, who require secondary prevention approaches, or to highly-selected patient subgroups, such as those with symptoms suggestive of CVD who require diagnostic strategies rather than risk assessment. Furthermore, these recommendations were not developed for use in specific subgroups of asymptomatic individuals at unusually high risk, such as those with genetically determined extreme values of traditional risk factors (e.g., patients with familial hypercholesterolemia).

2. Risk Assessment: Recommendations

Table 4. Summary of Recommendations for Risk Assessment

Recommendations	NHLBI Grade	NHLBI Evidence Statements	ACC/AHA COR	ACC/AHA LOE
1. The race- and sex-specific Pooled Cohort Equations* to predict 10-year risk for a first hard ASCVD event should be used in nonHispanic African Americans and nonHispanic Whites, 40 to 79 years of age.	B (Moderate)	N/A	I	B (4-8)

2. Use of the sex-specific Pooled Cohort Equations for nonHispanic Whites may be considered when estimating risk in patients from populations other than African Americans and nonHispanic Whites.	E (Expert Opinion)	Appendix 2 CQ2/ES1	IIb	C
3. If, after quantitative risk assessment, a risk-based treatment decision is uncertain, assessment of 1 or more of the following—family history, hs-CRP, CAC score, or ABI—may be considered to inform treatment decision making.	E (Expert Opinion)	Appendix 1	IIb†	B (9-17)
4. The contribution to risk assessment for a first ASCVD event using ApoB, CKD, albuminuria, or cardiorespiratory fitness is uncertain at present.	N (No Recommendation For or Against)	Appendix 1	N/A	N/A
5. CIMT is not recommended for routine measurement in clinical practice for risk assessment for a first ASCVD event.	N (No Recommendation For or Against)	Appendix 1	III: No Benefit†	B (12,16,18)
6. It is reasonable to assess traditional ASCVD risk factors‡ every 4 to 6 years in adults 20 to 79 years of age who are free from ASCVD and to estimate 10-year ASCVD risk every 4 to 6 years in adults 40 to 79 years of age without ASCVD.	B (Moderate)	Appendix 2 CQ2/ES7	IIa	B (19,20)
7. Assessing 30-year or lifetime ASCVD risk based on traditional risk factors‡ may be considered in adults 20 to 59 years of age without ASCVD and who are not at high short-term risk.	C (Weak)	Appendix 2 CQ2/ES2, CQ2/ES3, CQ2/ES4, CQ2/ES5, CQ2/ES6	IIb	C (20-22)

A downloadable spreadsheet enabling estimation of 10-year and lifetime risk for ASCVD and a web-based calculator are available at <http://my.americanheart.org/cvriskscalculator> and <http://www.cardiosource.org/science-and-quality/practice-guidelines-and-quality-standards/2013-prevention-guideline-tools.aspx>.

*Derived from the ARIC study (8), CHS (5), CARDIA study (23), Framingham original and offspring cohorts (4,6).

†Based on new evidence reviewed during ACC/AHA update of evidence.

‡Age, sex, total and HDL-cholesterol, systolic BP, use of antihypertensive therapy, diabetes, and current smoking.

ABI indicates ankle-brachial index; ACC, American College of Cardiology; AHA, American Heart Association; ApoB, Apolipoprotein B; ASCVD, atherosclerotic cardiovascular disease; BP, blood pressure; CAC, coronary artery calcium; CKD, chronic kidney disease; CIMT, carotid intima-media thickness; COR, Class of Recommendation; CQ, critical question, ES, evidence statement; HDL-C, high-density lipoprotein cholesterol; hs-CRP, high-sensitivity C-reactive protein; LOE, Level of Evidence; and NHLBI, National Heart, Lung, and Blood Institute.

3. Approach to Risk Assessment

In addressing its charge, the Work Group recognized the need for a risk assessment approach that was based on the types of data that primary care providers could easily collect and that could be implemented in routine clinical practice. After deliberation, the Work Group endorsed the existing and widely employed paradigm of matching the intensity of preventive efforts with the individual's absolute risk (24,25). The Work Group acknowledges that none of the risk assessment tools or novel risk markers examined in the present document have been formally evaluated in randomized controlled trials of

screening strategies with clinical events as outcomes. Nevertheless, this approach balances an understanding of an individual's absolute risk for CVD and potential treatment benefits against the potential absolute risks for harm from therapy. Using this framework, treatment can be targeted to those most likely to benefit without undue risk for harm, in the context of a "risk discussion." A risk discussion could include the assessment of the patient's risk for ASCVD, and potential benefits, negative aspects, risks, and patient preferences regarding initiation of relevant preventive therapies.

By its nature, such an approach requires a platform for reliable quantitative estimation of absolute risk based upon data from representative population samples. It is important to note that risk estimation is based on group averages that are then applied to individual patients in practice. This process is admittedly imperfect; no one has 10% or 20% of a heart attack during a 10-year period. Individuals with the same estimated risk will either have or not have the event of interest, and only those patients who are destined to have an event can have their event prevented by therapy. The criticism of the risk estimation approach to treatment-decision making also applies to the alternative, and much less efficient approach, of checking the patient's characteristics against numerous and complex inclusion and exclusion criteria for a potentially large number of pertinent trials. Only a small fraction of trial participants have events, and only a fraction of these events are prevented by therapy. Using either approach, the clinician must apply the average results obtained from groups of patients to the individual patient in practice.

Given the modification and adoption of the Framingham 10-year risk score for CHD risk assessment by the "Third Report of the National Cholesterol Education Program Expert Panel on Diagnosis, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III)" (25), and the uptake of this algorithm by practice sites across the United States, the Work Group began by discussing the value of retaining this algorithm. In collaboration with other NHLBI panels, the Work Group decided not to use this algorithm in its 2013 recommendations, because of its derivation in an exclusively White sample population and the limited scope of the outcome (in determining CHD alone). Rather, the Work Group derived risk equations from community-based cohorts that are broadly representative of the U.S. population of Whites and African Americans, and focused on estimation of first hard ASCVD events (defined as first occurrence of nonfatal myocardial infarction or CHD death, or fatal or nonfatal stroke) as the outcome of interest because it was deemed to be of greater relevance to both patients and providers. The focus on hard ASCVD, rather than CHD alone, is also consistent with evidence reviewed in a statement from the AHA/American Stroke Association calling for the inclusion of ischemic stroke in the outcome of interest for CVD risk assessment (26).

Numerous multivariable risk scores/equations have been derived and published (Appendix 3, and for more details, the Full Work Group Report Supplement (http://jaccjacc.cardiosource.com/acc_documents/2013_FPR_S5_Risk_Assessment.pdf). As part of its

deliberations, the Work Group considered previously published risk scores with validation in NHLBI cohort data as 1 possible approach. However, a number of persistent concerns with existing risk equations were identified including nonrepresentative or historically dated populations, limited ethnic diversity, narrowly defined endpoints, endpoints influenced by provider preferences (e.g., revascularizations), and endpoints with poor reliability (e.g., angina and heart failure [HF]). Given the inherent limitations of existing scores, the Work Group judged that a new risk score was needed to address some of the deficiencies of existing scores, such as utilizing a population sample that approaches, to the degree possible, the ideal sample for algorithm development and closely represents the U.S. population.

Data are sparse regarding usage and impact of absolute risk scores in clinical practice in primary prevention settings (27). Two systematic reviews, based on few studies, support the conclusion that risk assessment, combined with counseling, is associated with favorable but modest changes in patient knowledge and intention to change, and with provider prescribing behavior and risk factor control (28,29). No data are available on hard event outcomes. The Work Group specifically calls for research in this area (Section 8).

The Work Group notes that the “2009 ACCF/AHA Performance Measures for the Primary Prevention of CVD” specifically recommended use of global CVD risk estimation in clinical practice (30). Likewise, the U.S. Preventive Services Task Force recommendations for aspirin (31), NHLBI Adult Treatment Panel III recommendations (25), and European (32) and Canadian (33,34) guidelines for primary prevention of CVD, among others, have all recommended the use of absolute risk assessment for decision making about the intensity of lifestyle and pharmacological preventive interventions. Risk scores have been implemented in practice through paper scoring sheets, and increasingly through websites and downloadable applications. The electronic medical record can be adapted to estimate absolute risks automatically using patient data and published equations, and it is anticipated that risk estimation using this technology will become a mainstream application of the current and future risk algorithms.

4. Development of New Pooled Cohort ASCVD Risk Equations

Having made the decision to develop new equations to estimate the 10-year risk for developing a first ASCVD event, the Work Group used the best available data from community-based cohorts of adults, with adjudicated endpoints for CHD death, nonfatal myocardial infarction, and fatal or nonfatal stroke. Cohorts that included African-American or White participants with at least 12 years of follow-up were included. Data from other race/ethnic groups were insufficient, precluding their inclusion in the final analyses. The final pooled cohorts included participants from several large, racially and geographically diverse, modern NHLBI-sponsored cohort studies, including the ARIC (Atherosclerosis Risk in Communities) study (8), Cardiovascular Health Study (5), and the CARDIA (Coronary Artery Risk

Development in Young Adults) study (7), combined with applicable data from the Framingham Original and Offspring Study cohorts (4,6).

The Work Group used state-of-the-art statistical methods to derive and internally validate the Pooled Cohort Equations, which provide sex-and race-specific estimates of the 10-year risk for ASCVD for African-American and White men and women 40 to 79 years of age. The variables that statistically merit inclusion in the risk assessment equations are age, total and HDL-cholesterol, systolic BP (including treated or untreated status), diabetes, and current smoking status.

An expanded description of the derivation and validation of the Pooled Cohort Equations, as well as the means for implementing them in clinical practice, are provided in Appendix 4. Additional details are provided in the Full Report of the Work Group (http://jaccjacc.cardiosource.com/acc_documents/2013_FPR_S5_Risk_Assessment.pdf). A specific clinical vignette is also provided as an example in Appendix 4. In the clinical vignette, the 10-year risk is calculated for a patient 55 years of age who is a nonsmoker without diabetes, and with total cholesterol 213 mg/dL, HDL-cholesterol 50 mg/dL, and untreated systolic BP 120 mm Hg. Using these values in the Pooled Cohort Equations, the predicted 10-year ASCVD risks are 2.1% for White women, 3.0% for African-American women, 5.3% for White men, and 6.1% for African-American men.

Numerous other potential risk markers were considered for inclusion in the Pooled Cohort Equations, but for many there was no additional utility demonstrated upon their inclusion; for others, data were insufficient at the present time to determine their additional value. The equations were also assessed in external validation studies using data from other available cohorts. Other than the Framingham CHD risk score (and its derivative ATP-III risk assessment profile) and the European SCORE (System for Cardiac Operative Risk Evaluation) algorithm for CVD death, these equations have been subjected to more rigorous validation than other currently available equations, and they are the only risk assessment equations that include significant numbers of African Americans and focus on estimation of 10-year risk for the clinically relevant endpoint of ASCVD. The Work Group specifically calls for further research to develop similar equations applicable to other ethnic groups, to validate the utility of the Pooled Cohort Equations in diverse primary prevention settings, and to assess the potential benefit of novel risk markers when added to these equations, so that the equations may be modified or expanded over time as new data become available.

4.1. Recommendations for Assessment of 10-Year Risk for a First Hard ASCVD Event

Recommendation 1.

The race- and sex-specific Pooled Cohort Equations to predict 10-year risk for a first hard ASCVD* event should be used in nonHispanic African Americans and nonHispanic Whites, 40 to 79 years of age.

(Grade B, Moderate); ACC/AHA COR I, LOE B

Recommendation 2.

Use of the sex-specific Pooled Cohort Equations for nonHispanic Whites may be considered when estimating risk in patients from populations other than African Americans and nonHispanic Whites.

(Grade E, Expert Opinion); ACC/AHA COR IIb, LOE C

A downloadable spreadsheet enabling estimation of 10-year and lifetime risk for ASCVD and a web-based calculator are available at <http://my.americanheart.org/cvriskscalculator> and <http://www.cardiosource.org/science-and-quality/practice-guidelines-and-quality-standards/2013-prevention-guideline-tools.aspx>.

*Ten-year risk was defined as the risk of developing a first ASCVD event, defined as nonfatal myocardial infarction or CHD death, or fatal or nonfatal stroke, over a 10-year period among people free from ASCVD at the beginning of the period.

5. Implications for Risk Assessment

A range of estimated 10-year risk for a first hard ASCVD event is illustrated in the Full Work Group Report Supplement (Tables 8-11), across a broad range of risk factor burdens for selected combinations of the risk factors in sex-race groups (African-American and White women and men) (http://jaccjacc.cardiosource.com/acc_documents/2013_FPR_S5_Risk_Assesment.pdf). The estimated risks are specific to defined combinations of the risk factors, and demonstrate how they vary over a broad spectrum of potential profiles. Risk factor levels that are more adverse than those shown in these tables should always be associated with a higher estimated risk. For example, if a given risk factor combination indicates an estimated 10-year risk for hard ASCVD of 8%, but a patient has a higher level of systolic BP or total cholesterol, or a lower level of high-density lipoprotein cholesterol, than shown for that cell, then the estimated risk would be $\geq 8\%$. Because the estimated probabilities can become unstable when approaching the limits of the sample data, the risk probabilities are truncated at 1% and 30%. The proportion of the U.S. adult population, 40 to 79 years of age, in selected strata of estimated 10-year risk for hard ASCVD events, are shown overall and by sex and race in Table 5. When compared with nonHispanic Whites, estimated 10-year risk for ASCVD is generally lower in Hispanic-American and Asian-American populations and higher in American-Indian populations (35,36); hence, the lack of ethnic-specific risk algorithms are an important gap in our efforts to understand and prevent ASCVD in these populations. While the development of algorithms specific to these race/ethnic groups is encouraged, in the interim, providers may consider using the equations for nonHispanic Whites for these patients. When doing so, it is important to remember that the estimated risks may be over-estimates, especially for Hispanic- and Asian-Americans.

Table 5. Distribution of Estimated 10-Year Risk for a First Hard ASCVD Event in the CVD-Free, Nonpregnant U.S. Population 40 to 79 Years of Age, by Sex, and Race*

		Predicted 10-Year Risk for Hard ASCVD Event						
		<2.5%	2.5%–4.9%	5.0%–7.4%	7.5%–9.9%	10.0%–14.9%	15.0%–19.9%	≥20.0%
Total	% (95% CI) <i>n</i>	33.4 (31.2-35.5) 33,534,000	21.0 (19.4-22.7) 21,151,000	12.7 (11.4-14.0) 12,766,000	7.4 (6.5-8.3) 7,470,000	8.9 (8.1-9.6) 8,940,000	6.3 (5.6-7.1) 6,380,000	10.2 (9.5-11.0) 10,300,000
Sex								
Men	% (95% CI) <i>n</i>	17.4 (15.2-19.7) 8,386,000	22.7 (20.3-25.1) 10,950,000	15.6 (13.8-17.4) 7,511,000	10.1 (8.5-11.6) 4,847,000	12.1 (10.7-13.5) 5,849,000	8.8 (7.4-10.2) 4,248,000	13.3 (12.1-14.4) 6,388,000
Women	% (95% CI) <i>n</i>	48.0 (44.8-51.3) 25,148,000	19.5 (17.3-21.6) 10,200,000	10.0 (8.3-11.8) 5,256,000	5.0 (3.8-6.2) 2,622,000	5.9 (5.1-6.7) 3,091,000	4.1 (3.4-4.7) 2,131,000	7.5 (6.5-8.4) 3,912,000
Race								
White								
Men	% (95% CI) <i>n</i>	18.0 (15.0-21.1) 6,467,000	22.4 (19.4-25.3) 8,016,000	15.7 (13.3-18.1) 5,616,000	10.0 (8.2-11.8) 3,584,000	11.7 (9.9-13.5) 4,189,000	8.7 (7.0-10.4) 3,112,000	13.6 (12.3-14.9) 4,870,000
Women	% (95% CI) <i>n</i>	47.1 (43.0-51.1) 18,175,000	20.4 (17.7-23.0) 7,863,000	10.7 (8.6-12.8) 4,136,000	5.1 (3.6-6.7) 1,984,000	5.5 (4.6-6.5) 2,132,000	4.1 (3.4-4.9) 1,596,000	7.1 (5.9-8.2) 2,725,000
African American								
Men	% (95% CI) <i>n</i>	1.4 (0.3-2.6) 60,000	23.9 (19.9-28.0) 1,008,000	20.6 (17.0-24.2) 866,000	11.8 (8.8-14.8) 495,000	17.4 (14.3-20.5) 731,000	11.1 (8.2-13.9) 466,000	13.8 (11.0-16.7) 583,000
Women	% (95% CI) <i>n</i>	36.5 (32.4-40.6) 1,921,000	18.7 (15.6-21.8) 985,000	10.9 (8.6-13.2) 572,000	6.5 (5.0-7.9) 339,000	9.4 (7.2-11.7) 496,000	5.7 (4.2-7.2) 300,000	12.3 (9.5-15.0) 645,000
Hispanic								
Men	% (95% CI) <i>n</i>	24.0 (19.8-28.1) 1,303,000	22.1 (17.9-26.2) 1,200,000	13.2 (10.8-15.6) 718,000	10.6 (8.1-13.0) 574,000	11.4 (9.9-12.9) 619,000	6.2 (4.6-7.9) 339,000	12.6 (9.4-15.7) 683,000
Women	% (95% CI) <i>n</i>	59.4 (54.3-64.4) 3,293,000	14.5 (11.5-17.5) 803,000	7.5 (5.4-9.6) 418,000	4.5 (2.6-6.4) 248,000	4.9 (3.4-6.5) 273,000	3.0 (2.0-3.9) 164,000	6.3 (4.7-7.9) 347,000
Others								
Men	% (95% CI) <i>n</i>	20.8 (10.8-30.7) 555,000	27.1 (18.0-36.3) 726,000	11.6 (4.9-18.2) 310,000	7.2 (0.6-13.8) 193,000	11.5 (4.5-18.6) 309,000	12.3 (5.9-18.8) 330,000	9.4 (3.0-15.8) 251,000
Women	% (95% CI) <i>n</i>	59.8 (50.2-69.3) 1,757,000	18.6 (10.8-26.5) 548,000	4.4 (0-8.7) 128,000	1.7 (0-3.5) 49,000	6.4 (2.1-10.7) 188,000	2.4 (0.4-4.5) 71,000	6.7 (2.3-11.0) 195,000

Goff DC Jr, et al.
2013 ACC/AHA Cardiovascular Risk Guideline

*Data derived by applying the Pooled Cohort Equations to the National Health and Nutrition Examinations Surveys, 2007-2010 ($N=5,367$, weighted to 100,542,000 U.S. population)

ASCVD indicates atherosclerotic cardiovascular disease; CVD, cardiovascular disease; and U.S., United States.

6. CQs and Systematic Evidence Review

6.1. Critical Question 1

“What is the evidence regarding reclassification or contribution to risk assessment when hs-CRP, ApoB, GFR, microalbuminuria, family history, cardiorespiratory fitness, ABI, CAC, or CIMT are considered in addition to the variables that are in the traditional risk scores?”

The concept of matching the intensity of risk factor management to the estimated risk for CVD has been well established since the 27th Bethesda Conference in 1996 (24). As a consequence, widespread attention has focused on the accuracy and reliability of risk assessment. Claims that a minority of the risk for CVD can be explained by the major traditional risk factors, or that most patients presenting with CHD have no elevated traditional risk factors, have been disproven (37,38). Nonetheless, the desire to improve existing quantitative risk estimation tools has helped to stimulate and maintain interest in the search for new risk markers for CVD which might further enhance risk assessment.

CQ1 was developed to address whether newer risk markers have been identified that actually improve risk assessment enough to warrant routine measurement in clinical practice. This question applies to risk assessment in the general population, that is, the typical asymptomatic adult in routine clinical practice. This question does not address other highly selected patient subgroups, such as those with symptoms suggestive of CVD.

CQ1 was addressed using 2 independent approaches. First, in the process of developing the Pooled Cohort Equations, the additional risk markers listed in CQ1 were tested for inclusion in the model if they were available in the databases and could be evaluated on the basis of at least 10 years of follow up. A review of meta-analyses and systematic reviews published before September 19, 2013 was conducted in 2 stages. In the first stage, meta-analyses and systematic reviews published before April 2011 were identified and reviewed. In a second stage, conducted to update the evidence base before publication, additional meta-analyses and systematic reviews published before September 19, 2013 were identified and reviewed using the same criteria applied in the first stage. The reliance on published meta-analyses to evaluate novel biomarkers is a conservative approach that helps avoid the influence of positive publication bias that can occur early in the evaluation of a novel association and assures that we relied on a mature body of evidence (39).

Members of the Work Group proposed an initial list of novel risk markers for inclusion in CQ1 which was then prioritized during several rounds of discussion. In selecting the final list, the Work Group gave priority to factors that have engendered substantial discussion in the scientific community and that could be reasonably considered as potentially feasible for widespread population use by primary care providers in routine clinical settings in the United States. These deliberations considered availability, cost,

Goff DC Jr, et al.

2013 ACC/AHA Cardiovascular Risk Guideline

assay reliability, and risks of the test or downstream testing. The final list of new risk markers to be evaluated included several blood and urine biomarkers (hs-CRP, ApoB, creatinine [or estimated GFR], and microalbuminuria), several measures of subclinical CV disease (CAC, CIMT, and ABI), family history, and cardiorespiratory fitness. Other novel potential screening tools may be the subject of future guideline updates. When considering the utility of incorporating these new risk factors into routine risk assessment, guidance published by Hlatky et al (40) was considered. Special attention was given to the additional value these markers contributed to risk assessment in terms of discrimination, calibration, reclassification, and cost-effectiveness, in the context of any potential harm.

6.1.1. Summary of Systematic Reviews and Meta-Analyses for CQ1

Thirteen systematic review articles or meta-analyses met the inclusion/exclusion criteria (9-18,41-43). Publication dates ranged from 2008 to 2013. The Work Group reviewed the 13 systematic reviews and meta-analyses and created a table to list their key findings (Appendix 1). None of these markers has been evaluated as a screening test in randomized controlled trials with clinical events as outcomes. On the basis of current evidence, it is the opinion of the Work Group that assessments of family history of premature CVD, and measurement of hs-CRP, CAC, and ABI show some promise for clinical utility among the novel risk markers, based on limited data. Table 6 provides expert opinion regarding thresholds of these measures that may be considered for clinical decision making.

The Work Group notes that the review by Peters et al. (16) provides evidence to support the contention that assessing CAC is likely to be the most useful of the current approaches to improving risk assessment among individuals found to be at intermediate risk after formal risk assessment. Furthermore, the Work Group recognizes that the “2010 ACCF/AHA guideline for assessment of cardiovascular risk in asymptomatic adults” made recommendations regarding CAC testing (44). However, the Work Group notes that the outcomes in the studies reviewed by Peters et al. (16) and by Greenland et al. (44) were CHD outcomes, not hard ASCVD events that included stroke; hence, uncertainty remains regarding the contribution of assessing CAC to estimating 10-year risk of first hard ASCVD events after formal risk assessment using the new Pooled Cohort Equations. Furthermore, issues of cost and radiation exposure related to measuring CAC were discussed resulting in some uncertainty regarding potential risks of more widespread screening, which resulted in a decision in the current guideline to make assessment of CAC a Class IIb recommendation among individuals for whom a risk-based treatment decision is uncertain after formal risk estimation. The Work Group notes that this Class IIb recommendation is consistent with the recommendations in the 2010 ACCF/AHA guideline (44) for patients with a 10-year CHD risk of <10%, as well as for many other patients, because of the lower risk threshold (7.5% 10-year risk for a first hard ASCVD event) adopted by the “2013 ACC/AHA guideline on the treatment of blood cholesterol to

reduce atherosclerotic cardiovascular risk in adults” for recommending initiation of statin therapy for ASCVD risk reduction.

Furthermore, it was noted that measuring ApoB, albuminuria, GFR, or cardiorespiratory fitness is of uncertain value. Finally, the Work Group judged that the evidence provided by Den Ruijter et al (18) in combination with the concerns about measurement quality provided sufficient rationale to recommend against measuring CIMT in routine clinical practice for risk assessment for a first ASCVD event. If any of the 9 markers considered in this report is assessed in selected patients, the use of the information to guide treatment decisions will require sound clinician judgment and should be based on shared decision making.

Table 6. Expert Opinion Thresholds for use of Optional Screening Tests When Risk-Based Decisions Regarding Initiation of Pharmacological Therapy are Uncertain Following Quantitative Risk Assessment

Measure	Support Revising Risk Assessment Upward	Do Not Support Revising Risk Assessment
Family history of premature CVD	Male <55 years of age Female <65 years of age (1 st degree relative)	Occurrences at older ages only (if any)
hs-CRP	≥2 mg/L	<2 mg/L
CAC score	≥300 Agatston units or ≥75 th percentile for age, sex, and ethnicity*	<300 Agatston units and <75 percentile for age, sex, and ethnicity*
ABI	<0.9	≥0.9

*For additional information, see <http://www.mesa-nhlbi.org/CACReference.aspx>.

ABI indicates ankle-brachial index; CAC, coronary artery calcium; CVD, cardiovascular disease; and hs-CRP, high-sensitivity C-reactive protein.

6.1.2. Recommendations for CQ1: Use of Newer Risk Markers After Quantitative Risk Assessment

Recommendation 1. If, after quantitative risk assessment, a risk-based treatment decision is uncertain, assessment of 1 or more of the following—family history, hs-CRP, CAC score, or ABI—may be considered to inform treatment decision making.

(Grade E, Expert Opinion); ACC/AHA COR IIb, LOE B

Recommendation 2. CIMT is not recommended for routine measurement in clinical practice for risk assessment for a first ASCVD event.

(Grade N, No Recommendation For or Against); ACC/AHA Class III: No Benefit, LOE B

- *Based on new evidence reviewed during ACC/AHA update of the evidence.*

Recommendation 3. The contribution to risk assessment for a first ASCVD event using ApoB, chronic kidney disease, albuminuria, or cardiorespiratory fitness is uncertain at present.

(Grade N, No Recommendation For or Against)

6.2. Critical Question 2

“Are models constructed to assess the long-term (≥ 15 years or lifetime) risk for a first CVD event in adults effective in assessing variation in long-term risk among adults at low and/or intermediate short-term risk, whether analyzed separately or combined?”

A number of studies have noted that younger men (typically < 50 years of age) and most women have low (e.g., $< 5\%$ or $< 10\%$) predicted 10-year risks for CHD, and more broad CVD outcomes, despite the presence of significant risk factor burden (45,46). However, extensive epidemiological, pathological, and basic science data indicate that the development of atherosclerosis, the precursor of ASCVD, occurs over decades and is related to long-term and cumulative exposure to causal, modifiable risk factors. Thus, a life course perspective to risk assessment and prevention must be considered, especially among younger individuals. The primary value of risk factor measurement and quantitative long-term risk estimation in younger adults is 2-fold: first, to identify risk in individuals with extreme values of risk factors (e.g., familial hypercholesterolemia); second, to provide risk information and context regarding the potential benefits of lifestyle modification. When posing CQ2, the Work Group did not anticipate that long-term or lifetime risk would replace 10-year risk assessment as the foundation for absolute risk assessment and clinical decision-making. Rather, longer term risk estimates, if found to be useful, could provide adjunctive information for risk communication.

CQ2 was developed to assess the utility of long-term and lifetime risk assessment as an adjunct to short-term (10-year) risk assessment. It was recognized that there is little “disconnect” regarding approaches to prevention when the 10-year risk estimate is high (e.g., $> 10\%$ predicted 10-year risk): such patients merit intensive prevention efforts and should be considered for drug therapy to reduce or modify adverse levels of causal risk factors. CQ2 was selected for evaluation to determine whether quantitative or semi-quantitative long-term risk assessment would provide differential information that could be useful in risk communication, specifically to patients estimated to be at lower short-term risk. However, it is unclear what the long-term predicted and observed risks for CHD and CVD are among individuals who are at low predicted 10-year risk. CQ2 was designed to identify studies that assessed both short- and long-term risk, particularly focusing on those studies that provide long-term outcomes data for groups predicted to be at low 10-year risk. If a sufficiently large proportion of the population is at high long-term risk despite being at low short-term risk, then incorporating long-term risk assessment into routine clinical practice might have value for informing risk conversations with patients and guiding therapeutic lifestyle counseling and other aspects of care.

6.2.1. Summary of Evidence for CQ2

Ten studies that met inclusion/exclusion criteria were identified by the systematic review performed in April, 2011, and were examined (19-22,47-52). Publication dates ranged from 1999 to 2009. All of the

studies were observational. On the basis of these studies, 7 evidence statements were adopted (Appendix 2).

Multiple sources provided consistent evidence regarding the associations of traditional risk factors with events occurring during both short-term and long-term follow up. The important associations are best represented and understood in the context of multivariable risk equations that reliably predict absolute risk for ASCVD events. In addition, most of these risk factors are both causal and modifiable, indicating their central clinical importance for ASCVD prevention efforts. Given the additional evidence suggesting improved risk prediction using updated clinical covariates, the Work Group makes the following recommendations.

6.2.2. Recommendations for CQ2: Long-Term Risk Assessment

Recommendation 1. It is reasonable to assess traditional ASCVD risk factors every 4 to 6 years in adults 20 to 79 year of age who are free from ASCVD and estimate 10-year ASCVD risk every 4 to 6 years in adults 40 to 79 years of age who are free from ASCVD.

(Grade B, Moderate); ACC/AHA COR IIa, LOE B

Recommendation 2. Assessing 30-year or lifetime ASCVD risk based on traditional risk factors[†] may be considered in adults 20 to 59 years of age who are free from ASCVD and who are not at high short-term risk.

(Grade C, Weak); ACC/AHA COR IIb, LOE C

A downloadable spreadsheet enabling estimation of 10-year and lifetime risk for ASCVD and a web-based calculator are available at <http://my.americanheart.org/cvriskscalculator> and <http://www.cardiosource.org/science-and-quality/practice-guidelines-and-quality-standards/2013-prevention-guideline-tools.aspx>.

[†]Age, sex, total and HDL-cholesterol, systolic BP, use of antihypertensive therapy, diabetes, and current smoking.

Evidence was not found regarding the utility of lifetime risk assessment for guiding pharmacologic therapy decisions, and the Work Group judged that long-term and lifetime risk information may be used more appropriately at this time to motivate therapeutic lifestyle change in younger individuals. This perspective influenced the choice of age 20 as the starting point for long-term risk assessment, despite a threshold of age 40 for short-term 10-year ASCVD risk assessment.


Long-term and lifetime risk estimation may be less valuable for individuals who are found to be at high short-term (10-year) risk based on multivariable equations in whom decisions regarding prevention efforts may be clear. However, an understanding of long-term risk may provide a means for encouraging adherence to lifestyle or pharmacological therapies, especially for patients who might have difficulty understanding the importance of their short-term risk. Likewise, for older individuals, or those

with limited life expectancy, clinical considerations should dictate the intensity of risk assessment and prevention efforts.

7. Implementation Considerations for Risk Assessment

A suggested approach for incorporating these recommendations into clinical practice is shown in Figure 1. For patients 20 to 79 years of age who are free from clinical ASCVD, the first step is to assess ASCVD risk factors. Whereas it is reasonable to assess ASCVD risk factors in younger and older individuals, limitations in available data prevented the development of robust risk assessment algorithms in these populations. Hence, for patients outside this age range, providers should refer to applicable clinical practice guidelines (i.e., pediatric (53) and adult primary prevention guidelines (54,55)). Risk assessment should be repeated every 4 to 6 years in persons who are found to be at low 10-year risk ($<7.5\%$). Beginning at age 40, formal estimation of the absolute 10-year risk for ASCVD is recommended. Long-term or lifetime risk estimation is recommended for all persons who are between 20 to 39 years of age and for those between 40 to 59 years of age who are determined to be at low 10-year risk ($<7.5\%$). As shown in Figure 1, all patients should receive applicable risk information and appropriate lifestyle counseling. The 10-year risk estimates provided by the new Pooled Cohort Equations differ from those generated by the Adult Treatment Panel III algorithm in several respects (25) as discussed in detail in the Full Work Group Report (http://jaccjacc.cardiosource.com/acc_documents/2013_FPR_S5_Risk_Assesment.pdf). To summarize, based on the risk estimation algorithm recommended by Adult Treatment Panel III, approximately 31.9% of the ASCVD-free, nonpregnant U.S. population between 40 and 79 years of age have a 10-year risk of a first hard CHD event of at least 10% or have diabetes. Based on the new Pooled Cohort Equations described here, approximately 32.9% have a 10-year risk of a first hard ASCVD of at least 7.5%. The outcomes and thresholds of these 2 approaches are different, but the overlap of these 2 means for defining high-risk groups is substantial, at roughly 75%. Nonetheless, these important differences make simple linear conversions imprecise. We recommend that healthcare organizations convert to these new Pooled Cohort Equations as soon as practical (Appendix 4). A downloadable spreadsheet enabling estimation of 10-year and lifetime risk for ASCVD and a web-based calculator are available at <http://my.americanheart.org/cvriskcalculator> and <http://www.cardiosource.org/science-and-quality/practice-guidelines-and-quality-standards/2013-prevention-guideline-tools.aspx>.

Figure 1. Implementation of Risk Assessment Work Group Recommendations



ACC indicates American College of Cardiology; AHA, American Heart Association; and ASCVD, atherosclerotic cardiovascular disease.

8. Evidence Gaps and Future Research Needs

The Work Group strongly recommends continued research to fill gaps in knowledge regarding short- and long-term ASCVD risk assessment and outcomes in all race/ethnic groups, across the age spectrum, and in women and men. Future research should include analyses of short- and long-term risk in diverse groups; optimal communication of ASCVD risk information; utility of short-and long-term risk assessment for motivating behavioral change and adherence to therapy; utility of short-and long-term risk assessment for influencing risk factor levels and clinical outcomes; utility of differential information conveyed by short- and long-term risk assessment; and utility of novel risk markers in short- and long-term risk assessment.

9. Conclusions

The Work Group's approach to risk assessment represents a step forward in ASCVD prevention that is large enough to justify the challenges inherent in implementing a new approach, rather than staying with the CHD risk assessment approach recommended previously. The final recommendations are summarized in Table 4 and Figure 1. Two major advantages of this approach are the ability to estimate risk for a broader based ASCVD outcome that is more relevant to additional segments of the population, including women and African Americans, and the ability to provide risk estimates specific to African Americans.

Goff DC Jr, et al.
2013 ACC/AHA Cardiovascular Risk Guideline

Promoting lifetime risk estimation may represent an additional step forward in supporting lifestyle behavior change counseling efforts. Periodic updating of the guidelines should address numerous issues related to risk assessment.

Presidents and Staff

American College of Cardiology Foundation

John Gordon Harold, MD, MACC, President
 Shalom Jacobovitz, Chief Executive Officer
 William J. Oetgen, MD, MBA, FACC, Executive Vice President, Science, Education & Quality
 Charlene May, Senior Director, Science and Clinical Policy

American College of Cardiology Foundation/American Heart Association

Lisa Bradfield, CAE, Director, Science and Clinical Policy

American Heart Association

Mariell Jessup, MD, FACC, FAHA, President
 Nancy Brown, Chief Executive Officer
 Rose Marie Robertson, MD, FAHA, Chief Science Officer
 Gayle R. Whitman, PhD, RN, FAHA, FAAN, Senior Vice President, Office of Science Operations
 Marco Di Buono, PhD, Vice President of Science and Research
 Jody Hundley, Production Manager, Scientific Publications, Office of Science Operations

National Heart, Lung, and Blood Institute

Denise Simons-Morton, MD, PhD

RTI International

Harmon Jordan, ScD

Science Applications International Corporation

Lev Nevo, MD

Acknowledgment: The Work Group would like to acknowledge the significant contributions of Dr. Hongyan Ning, Biostatistician at Northwestern University Feinberg School of Medicine, in providing statistical analyses for this document.

Key Words: TBD

Appendix 1. Evidence Statements for CQ1

ES Number	Author/Group	Factor	Evidence Statement/Conclusion
1	USPSTF (9)	hs-CRP	<p>“Strong evidence indicates that CRP is associated with CHD events. Moderate, consistent evidence suggests that adding CRP to risk prediction models among initially intermediate-risk persons improves risk stratification.”</p> <p>“Few studies directly assessed the effect of CRP on risk reclassification in intermediate-risk persons.”</p> <p>hs-CRP was associated with risk and results in some reclassification in intermediate-risk persons, but it was not clear whether this reclassification led to a net improvement in prediction. Values of receiver operating curve C-statistics, measures of discrimination, are mentioned but not reported; hence, no evidence on discrimination, calibration, net reclassification index or cost-effectiveness was provided. Reports some impact on reclassification, probably modest (pp. 488–491).</p>
2	Helfand et al., 2009 (12)	hs-CRP, CAC, CINT, ABI	<p>With respect to risk assessment for major CHD, the authors concluded that, “The current evidence does not support the routine use of any of the 9 risk factors for further risk stratification of intermediate-risk persons.” The nine risk factors examined were: hs-CRP, CAC score as measured by electron-beam computed tomography, lipoprotein (a) level, homocysteine level, leukocyte count, fasting blood glucose, periodontal disease, ABI, and CINT.</p> <p>hs-CRP was associated with CHD and led to some reclassification. The authors cite the JUPITER results to support the conclusion that hs-CRP testing may be useful in intermediate-risk patients to drive statin therapy. The Work Group recognizes that more recent individual study results have been published. Updated systematic reviews addressing discrimination, calibration, reclassification, and cost issues in the context of the newer ASCVD risk assessment model proposed in this document are needed.</p> <p>CAC was associated with CHD and with some reclassification, but it is uncertain how much and how valuable this reclassification is. The document provides little evidence regarding discrimination, calibration, and cost-effectiveness. The Work Group also is concerned about radiation and incidental findings. The Work Group recognizes that more recent individual study results have been published. Updated systematic reviews addressing discrimination, calibration, reclassification, cost, and safety issues in the context of the newer ASCVD risk assessment model proposed in this document are needed.</p> <p>CINT was associated with CHD, but the document provides little evidence regarding reclassification, discrimination, calibration, and cost-effectiveness. The Work Group also has concerns about measurement issues. Standardization of CINT measurement is a major challenge. The Work Group recognizes that more recent individual study results have been published. Updated systematic reviews addressing discrimination, calibration, reclassification, cost, and measurement (standardization) issues in the context of the newer ASCVD risk assessment model proposed in this document are needed.</p>

Goff DC Jr, et al.
2013 ACC/AHA Cardiovascular Risk Guideline

ES Number	Author/Group	Factor	Evidence Statement/Conclusion
			ABI was associated with CHD and some reclassification, but it is uncertain how much and how valuable this reclassification is. Evidence suggests some improvement in discrimination, but the document provides little evidence regarding calibration and cost-effectiveness. The Work Group members are uncertain whether more recent individual study results have been published relevant to ABI. Updated systematic reviews addressing discrimination, calibration, reclassification, and cost issues in the context of the newer ASCVD risk assessment model proposed in this document are needed.
3	Emerging Risk Factors Collaboration (13)	hs-CRP	<p>“CRP concentration has continuous associations with the risk for coronary heart disease, ischaemic stroke, vascular mortality, and death from several cancers and lung disease that are each of broadly similar size. The relevance of CRP to such a range of disorders is unclear. Associations with ischaemic vascular disease depend considerably on conventional risk factors and other markers of inflammation.”</p> <p>hs-CRP is associated with risk for CVD. This analysis did not directly assess value in risk prediction. No additional evidence was provided regarding discrimination, calibration, reclassification, or cost-effectiveness.</p>
4	Schnell-Inderst et al., 2010 (17)	hs-CRP	<p>For MI and cardiovascular mortality, “Adding hs-CRP to traditional risk factors improves risk prediction, but the clinical relevance and cost-effectiveness of this improvement remain unclear.”</p> <p>Absolute differences in C-statistics between models including and not including hs-CRP ranged from 0.00 to 0.027.</p> <p>Some evidence was provided to support the cost-effectiveness of hs-CRP testing in some modeling scenarios, characterized by intermediate- and higher-risk populations and lower cost (generics) statins of at least moderate efficacy.</p>
5	Emerging Risk Factors Collaboration (41)	ApoB	This paper provided evidence of rough equivalence of associations of CVD with non-HDL-C and ApoB after multivariable adjustment (including HDL-C). See Figure 1 for CHD and the text for stroke. By inference, this finding means there would be rough equivalence between ApoB and total cholesterol with similar adjustment.
6	Sniderman et al., 2011 (43)	ApoB	ApoB was more strongly related to risk for ASCVD than either non-HDL-C or LDL-C in a substitution model that also included HDL-C. No evidence was presented pertinent to an addition model in which ApoB might be added to a model that included total cholesterol, LDL-C or non-HDL-C. Additional models are the type of model of interest to this question. By inference, these results may mean that ApoB

Goff DC Jr, et al.
2013 ACC/AHA Cardiovascular Risk Guideline

ES Number	Author/Group	Factor	Evidence Statement/Conclusion
			is more strongly related to risk than is total cholesterol. This paper did not address directly the value of adding ApoB to a model with traditional risk factors. No information was presented regarding discrimination, calibration, reclassification, or cost. The relative risks evaluated in the meta-analysis were adjusted for various sets of covariates in the various primary reports, and the adjustments were judged to be incomplete. Furthermore, studies of varying designs and quality were included, leaving the Work Group members concerned regarding the validity of the evidence.
7	Kodama et al., 2009 (42)	Cardiorespiratory fitness	Better cardiorespiratory fitness was associated with lower risk for all-cause mortality and CHD/CVD. Based on the sensitivity analyses in table 2, evidence of association was weaker for CHD/CVD, but still significant, when based on studies with more complete adjustment for other risk factors. The utility of assessing cardiorespiratory fitness in risk prediction was not assessed (discrimination, calibration, reclassification and cost).
8	Ankle Brachial Index Collaboration (11)	ABI	ABI is associated with total CHD risk and leads to significant reclassification, and the pattern of reclassification is different by sex. Among men, the effect is to down-classify high-risk men. Among women the effect is to up-classify low-risk women. Overall, the FRS, as applied by the investigators, showed relatively poor discrimination in this meta-analysis, with <i>C</i> -statistics of 0.646 (95% CI: 0.643–0.657) in men and 0.605 (0.590–0.619) in women. There was an improvement in <i>C</i> -statistic in both men, 0.655 (0.643–0.666) and women 0.658 (0.644–0.672) when ABI was added to a model with FRS. The improvement in the <i>C</i> -statistic was greater and significant in women but was not significant in men. No evidence on calibration, net reclassification index, or cost-effectiveness was provided.
9	Empana, et al, 2011 (10)	Family history of CHD	<p>“In separate models adjusted for age, gender, and study cohort, a family history of CHD, BMI, and waist circumference were all predictors of CHD. When traditional risk factors were controlled for, family history of CHD ($p<0.001$) and BMI ($p=0.03$) but not waist circumference ($p=0.42$) remained associated with CHD. However, the addition of family history of CHD or BMI to the traditional risk factors model did not improve the discrimination of the model (not shown).”</p> <p>This paper developed a CHD risk prediction algorithm based on 4 French population studies, and evaluated, among other factors, the contribution of family history to traditional risk factors. Family history of CHD was defined as the self-report of a myocardial infarction (MI) in first degree relatives (parents and siblings) in the D.E.S.I.R. and SU.VI.MAX studies, as a history of MI before 55 years in men and before 65 years in women in parents, siblings, and grandparents in the PRIME study, and as a death due to MI in</p>

Goff DC Jr, et al.
2013 ACC/AHA Cardiovascular Risk Guideline

ES Number	Author/Group	Factor	Evidence Statement/Conclusion
			first degree relatives in the Three City study. No evidence on calibration, net reclassification index, or cost-effectiveness was provided.
10	Moyer et al. 2013 (15)	ABI	<p>This paper is an updated review of the utility of assessing ABI for the USPSTF.</p> <p>“The USPSTF concludes that the current evidence is insufficient to assess the balance of benefits and harms of screening for PAD and CVD risk assessment with the ABI in adults. (I statement)”</p> <p>“The USPSTF found no evidence that screening for and treatment of PAD in asymptomatic patients leads to clinically important benefits. It also reviewed the potential benefits of adding the ABI to the FRS and found evidence that this results in some patient risk reclassification; however, how often the reclassification is appropriate or whether it results in improved clinical outcomes is not known.”</p> <p>The Work Group notes that this review provides some evidence that assessing ABI may improve risk assessment; however, no evidence was found by the USPSTF reviewers pertinent to the question of whether measuring ABI leads to better patient outcomes.</p>
11.	Peters et al. 2012 (16)	CIMT, CAC	<p>This paper is a systematic review of the literature regarding the contribution to risk assessment of imaging for subclinical atherosclerosis.</p> <p>“Published evidence on the added value of atherosclerosis imaging varies across the different markers, with limited evidence for FMD and considerable evidence for CIMT, carotid plaque and CAC. The added predictive value of additional screening may be primarily found in asymptomatic individuals at intermediate cardiovascular risk. Additional research in asymptomatic individuals is needed to quantify the cost effectiveness and impact of imaging for subclinical atherosclerosis on cardiovascular risk factor management and patient outcomes.”</p> <p>Regarding CIMT:</p> <p>“The c-statistic of the prediction models without CIMT increased from 0.00 to 0.03 when CIMT was added. In the Atherosclerosis Risk In Communities (ARIC) study, addition of CIMT to the prediction model resulted in an NRI overall of 7.1% (95% CI 2.2% to 10.6%) and an IDI of 0.007 (95% CI 0.004 to 0.010). The NRI intermediate was 16.7% (95% CI 9.3% to 22.4%). In contrast, 10 year results from the Carotid Atherosclerosis Progression Study showed that addition of CIMT to the prediction model resulted in an IDI of 0.04% and NRI overall of -1.41%. Analysis of 1574 participants from the Firefighters and Their Endothelium study showed an NRI overall of 11.6% (p=0.044) and an NRI intermediate of 18.0%</p>

Goff DC Jr, et al.
2013 ACC/AHA Cardiovascular Risk Guideline

ES Number	Author/Group	Factor	Evidence Statement/Conclusion
			<p>(p=0.034).”</p> <p>The Work Group notes that this paper provides some evidence to consider assessing CIMT; however, this conclusion was not supported by the Den Ruijter article described below.</p> <p>Regarding CAC:</p> <p>“The c-statistic increased from 0.04 to 0.13 when CAC was added to the model. Four recently published studies also reported results on the NRI and/or the IDI. One of these studies comprised a subgroup analysis of an earlier publication in the total population in individuals without indications for statin therapy. Analyses of the MESA study showed that addition of CAC to the conventional prediction model resulted in an NRI overall of 25% (95% CI 16% to 34%) and an NRI intermediate of 55% (95% CI 41% to 69%). The IDI in the MESA study was 0.026. Results were similar in the Rotterdam study. Addition of CAC to the prediction model led to an NRI overall of 14% (p<0.01) which was mainly driven by correctly reclassifying those at intermediate risk according to the traditional prediction model. Results from the Heinz Nixdorf Recall study also showed large NRIs when CAC was added to the Framingham Risk Score. Using different thresholds to define the intermediate risk category (10-20% or 6-20%), the NRI overall was 22% and 20%, respectively. The NRI intermediate was 22% for intermediate risk thresholds of 10-20% and 31% for intermediate risk thresholds of 6-20%. In addition, the IDI was 0.0152 when the prediction models with and without CAC were compared. The NRI overall was 25.1% and the IDI was 0.0167 in individuals from the Heinz Nixdorf Recall study without indications for statin therapy.”</p> <p>The Work Group notes that this paper provides evidence to support the conclusion that assessing CAC is likely to be the most useful approach to improving risk assessment among individuals found to be at intermediate risk after formal risk assessment. Furthermore, we note that the outcomes in the studies reviewed above were CHD, not ASCVD. The Work Group discussed concerns about cost, radiation exposure and the uncertainty of the contribution of assessing CAC to estimating 10-year risk of hard ASCVD after formal risk assessment.</p>
12.	Kashani et al, 2013 (14)	Family history	<p>This paper is an integrative literature review on the contribution of assessing family history to risk appraisal.</p> <p>“The evidence demonstrates that family history is an independent contributor to risk appraisal and unequivocally supports its incorporation to improve accuracy in global CVD risk estimation.”</p>

Goff DC Jr, et al.
2013 ACC/AHA Cardiovascular Risk Guideline

ES Number	Author/Group	Factor	Evidence Statement/Conclusion
			The Work Group notes that a variety of endpoints, clinical and subclinical, were included in the reviewed papers. No evidence on discrimination, calibration, net reclassification index, or cost-effectiveness was provided.
13.	Den Ruijter et al, 2012 (18)	CIMT	<p>This paper is an individual level meta-analysis of “14 population-based cohorts contributing data for 45 828 individuals. During a median follow-up of 11 years, 4007 first-time myocardial infarctions or strokes occurred.”</p> <p>“We first refitted the risk factors of the FRS and then extended the model with common CIMT measurements to estimate the absolute 10-year risks to develop a first-time myocardial infarction or stroke in both models. The C statistic of both models was similar (0.757; 95% CI, 0.749-0.764; and 0.759; 95% CI, 0.752-0.766). The net reclassification improvement with the addition of common CIMT was small (0.8%; 95% CI, 0.1%-1.6%). In those at intermediate risk, the net reclassification improvement was 3.6% in all individuals (95% CI, 2.7%-4.6%) and no differences between men and women.”</p> <p>“The addition of common CIMT measurements to the FRS was associated with small improvement in 10-year risk prediction of first-time myocardial infarction or stroke, but this improvement is unlikely to be of clinical importance.”</p> <p>The Work Group judged this paper to provide the strongest evidence available regarding the potential value of CIMT to risk assessment. The Work Group also has concerns about measurement issues. Standardization of CIMT measurement is a major challenge.</p>

ABI indicates ankle-brachial index; ApoB, apolipoprotein B; BMI, body mass index; ASCVD, atherosclerotic cardiovascular disease; CVD, cardiovascular disease; CAC, coronary artery calcium; CHD, coronary heart disease; CIMT, carotid intima-media thickness; ES, evidence statement; FRS, Framingham Risk Score; HDL-C, high-density lipoprotein cholesterol; hs-CRP, high-sensitivity C-reactive protein; JUPITER, Justification for the Use of Statins in Primary Prevention: An Intervention Trial Evaluating Rosuvastatin; LDL-C, low-density lipoprotein cholesterol; MI, myocardial infarction; and USPSTF, United States Preventive Services Task Force.

Appendix 2. Evidence Statements for CQ2

Evidence Statement	References
<p>1. We found no evidence assessing variations in long-term or lifetime risk for CVD outcomes among persons at low or intermediate short-term risk in race/ethnic groups other than nonHispanic Whites in the United States and Europe.</p> <p><i>Strength of Evidence: None</i></p>	--
<p>2. ASCVD risk factors measured in young and middle-aged adults, considered singly or jointly, generally are associated with short-term (≤ 10 years), long-term (≥ 15 years), and lifetime risk for ASCVD.</p> <p><i>Strength of Evidence:</i> Low (for diabetes and metabolic syndrome) to Moderate (for BMI, cholesterol, systolic BP, and smoking).</p>	(20,21,47,48,51,52)
<p>3. Multivariable short-term (10-year) CHD risk prediction models underestimate absolute lifetime risk for CHD, but may stratify relative lifetime risk for CHD in women and older men.*</p> <p><i>Strength of Evidence: Low</i></p> <p>*CHD is defined as all manifestations of CHD, or as CHD death/nonfatal MI.</p>	(22)
<p>4. Long-term (30-year) risk equations based on traditional ASCVD risk factors* provide more accurate prediction of long-term ASCVD† risk than do extrapolations of short-term (10-year) risk equations among individuals 20 to 59 years of age free from ASCVD.</p> <p><i>Strength of Evidence: Low</i></p> <p>*Age, sex, total and HDL-C, systolic BP, use of antihypertensive therapy, diabetes, current smoking</p> <p>†CHD death, nonfatal MI, or fatal/nonfatal stroke; or all ASCVD</p>	(20)
<p>5. The presence and severity of selected traditional ASCVD risk factors* stratify absolute levels of lifetime risk for ASCVD† among nonHispanic White adults 45 to 50 years of age who are free of ASCVD and not at high short-term risk.</p> <p><i>Strength of Evidence: Low</i></p> <p>*Risk factors were considered in 5 mutually exclusive strata encompassing the full spectrum of risk levels, as follows: 1) ≥ 2 major</p>	(21)

Goff DC Jr, et al.
2013 ACC/AHA Cardiovascular Risk Guideline

<p>risk factors (defined as total cholesterol ≥ 240 mg/dL or treated, systolic BP ≥ 160 or diastolic BP ≥ 100 mm Hg or treated, or diabetes, or current smoking), lifetime risk for ASCVD $>50\%$; 2) 1 major risk factor only, lifetime risk for ASCVD 39% to 50%; 3) ≥ 1 elevated risk factors (defined as untreated total cholesterol 200 to 239 mg/dL, or untreated systolic BP 140 to 159 mm Hg or diastolic BP 90 to 99 mm Hg, and no diabetes and no current smoking), lifetime risk for ASCVD 39% to 46%; 4) 1 or more risk factors at nonoptimal levels (untreated total cholesterol 180 to 199 mg/dL, or untreated systolic BP 120 to 139 mm Hg or diastolic BP 80 to 89 mm Hg, and no diabetes and no current smoking), lifetime risk for ASCVD 27% to 36%; and 5) all optimal levels of risk factors (defined as untreated total cholesterol <180 mg/dL, and untreated BP <120/<80 mm Hg, and no diabetes, and no current smoking), lifetimes risk for ASCVD $<10\%$.</p> <p>†CHD death, MI, coronary insufficiency, angina, fatal/nonfatal atherothrombotic stroke, claudication, other CVD death</p>	
<p>6. Long-term (≥ 15 years) risk prediction models based on selected traditional ASCVD risk factors* predict CHD death with good discrimination and calibration, and better in women than men, in U.S. nonHispanic White populations.</p> <p>Strength of Evidence: Low</p> <p>*Age, sex, total cholesterol, systolic BP, diabetes, smoking</p>	(50)
<p>7. Measuring and updating ASCVD risk factors every 4 to 6 years improves short- and long-term risk prediction.</p> <p>Strength of Evidence: Moderate</p>	(19,20)

ASCVD indicates atherosclerotic cardiovascular disease; BMI, body mass index; BP, blood pressure; CHD, coronary heart disease; CVD, cardiovascular disease; CQ, critical question; HDL-C, high-density lipoprotein cholesterol; IDI, improvement index; MI, myocardial infarction; NRI, net reclassification index; PAD, peripheral artery disease; and --, none.

**Appendix 3. Characteristics of Previously Published Risk Scores and Current Pooled Cohort Equations
(Including Data Sources, Covariates, and Outcomes)**

Risk Score				Risk Factors/Covariates Included															Cardiovascular Disease Events								
																			Hard CVD including cardiac failure			Hard ASCVD			Hard CHD		
																			Total CHD								
																			Total CHD including revascularization								
																			Coronary Revasc	Angina Pectoris	Unstable Angina	Myocardial Infarct	CHD Death	Stroke	Stroke Death	Cardiac Failure	TIA
Study Group	Study and Region	Data Source	Publication Year	Age	Sex	Total Chol	LDL-Chol	HDL-Chol	CRP	Systolic BP	BP Rx	Diabetes	HbA1c*	Smoking	Family Hx CVD†	Body Mass Index	Social	Region	Coronary Revasc	Angina Pectoris	Unstable Angina	Myocardial Infarct	CHD Death	Stroke	Stroke Death	Cardiac Failure	TIA
Framingham CHD (56)	Framingham MA, USA	EAF, EAM	1998	x	x	x	x	X		x		x		x						x	x	X	x				
ATP III (25)	Framingham MA, USA	EAF, EAM	2001	x	x	x		X		x	x			x								X	x				
Framingham Global (57)	Framingham MA, USA	EAF, EAM	2008	x	x	x		X		x	x	x		x								X	x	x	x	x	
PRO-CAM (58)	Muenster, Germany	EM	2002	x			x	X		x		x		x	x							X	x				
QRISK (59)	QRESEARCH, United Kingdom	EF, EM	2007	x	x	x		X		x	x			x	x	x	x [‡]	x	x	x	x	X	x	x	x		x
Reynolds Men (60)	Phys Health Study USA	EAF	2008	x		x		X	x	x				x	x				x			X	x	x	x		
Reynolds Women (61)	Women's Health Study USA	EAM	2007	x		x		X	x	x			x	x	x				x			X	x	x	x		
EUROSCORE (62)	12 cohorts Europe	EF, EM	2003	x	x	x		X		x				x				x					x		x		

Goff DC Jr, et al.
2013 ACC/AHA Cardiovascular Risk Guideline

Risk Score				Risk Factors/Covariates Included															Cardiovascular Disease Events									
																						Hard CVD including cardiac failure						
																						Hard ASCVD						
																						Hard CHD						
																				Total CHD								
																			Total CHD including revascularization									
Study Group	Study and Region	Data Source	Publication Year	Age	Sex	Total Chol	LDL-Chol	HDL-Chol	CRP	Systolic BP	BP Rx	Dia-betes	HbA1c*	Smok-ing	Family Hx CVD†	Body Mass Index	Social	Region	Coron-ary Revasc	Angina Pectoris	Unsta-ble Angina	Myo-cardial Infarct	CHD Death	Stroke	Stroke Death	Cardiac Failure	TIA	
Pooled Cohort (current)	CARDIA, Framingham, ARIC, CHS,USA	EAF, EAM, AAF, AAM		x	x	x		X		x	x	x		x								X	x	x	x			

Risk calculators noted above include hyperlinks to the respective webpage.

* Only among those with diabetes

† Definitions of a positive family history vary

‡ Measure of social deprivation

AAF indicates African-American females; AAM, African-American males; ASCVD, atherosclerotic cardiovascular disease; BP, blood pressure; Chol, cholesterol; CHD, coronary heart disease; CRP, C-reactive protein; CVD, cardiovascular disease; EF, European females; EM, European males; EAF, European American females; EAM, European American males; HbA1c, hemoglobin A1c; Hx, history; Revasc, revascularization; and TIA, transient ischemic attack.

Appendix 4. Development and Steps for Implementation of the ASCVD Pooled Cohort Risk Equations

Prior experience with the development of the Framingham Heart Study 10-year *CHD* risk prediction equations (25,56), and the more recent Framingham 10-year *general CVD* risk prediction equations (63), were used as a basis for developing the new Pooled Cohort Risk Equations. To expand the utility and generalizability of the new equations, extensive data were used from several large, racially and geographically diverse, modern NHLBI-sponsored cohort studies, including the ARIC (Atherosclerosis Risk in Communities) study (8), Cardiovascular Health Study (5), and the CARDIA (Coronary Artery Risk Development in Young Adults) study (7), combined with applicable data from the Framingham Original and Offspring Study cohorts (4,6).

A total of 11,240 White women (who experienced 902 hard ASCVD events), 9,098 White men (1,259 hard ASCVD events), 2,641 African-American women (290 hard ASCVD events), and 1,647 African-American men (238 hard ASCVD events) who met the following criteria were included: 40 to 79 years of age, apparently healthy, and free of a previous history of nonfatal myocardial infarction (recognized or unrecognized), stroke, HF, percutaneous coronary intervention, coronary artery bypass surgery, or atrial fibrillation. Data from the included participants were used to develop sex- and race-specific equations to predict 10-year risk for a first hard ASCVD event. Due to the growing health burden of HF, the Work Group examined the possibility of including HF as an outcome. However, study-by-study ascertainment and adjudication of HF varied considerably, and therefore HF could not be included as an outcome. Due to known substantial geographic variation in use, (Dartmouth Atlas of Healthcare, <http://www.dartmouthatlas.org/>) self-selection, and physician recommendation biases (64), coronary revascularization was also not included as an endpoint.

The Pooled Cohort Equations for estimating ASCVD were developed from sex- and race-specific proportional hazards models that included the covariates of age, treated or untreated systolic BP level, total cholesterol and high-density lipoprotein cholesterol levels, current smoking status (Y/N), and history of diabetes (Y/N). A variable representing lipid treatment was considered, but not retained in the final model because lipid therapy was relatively uncommon in the cohorts and statistical significance was lacking. Baseline characteristics of the participants included in the equation derivation model are shown in the Full Panel Report Data Supplement, as are details of the methods used to derive, evaluate, and validate (internally and externally) the resulting risk equations and their potential limitations. In summary, discrimination and calibration of the models were very good. C statistics ranged from a low of 0.713 (African-American men) to a high of 0.818 (African-American women). Calibration chi-square statistics ranged from a low of 4.86 (nonHispanic White men) to a high of 7.25 (African-American women). The coefficients for the equations for calculating an estimate of an individual's 10-year risk for a first hard

ASCVD event are provided in Table A, along with examples based on a specific risk profile for each race-sex group. The step-by-step process for estimating the risk in the specific examples of Table A is provided in Table B. These 2 tables are intended to enable programmers to integrate these equations into electronic health records.

Table A. Equation Parameters of the Pooled Cohort Equations for Estimation of 10-Year Risk for Hard ASCVD* and Specific Examples for Each Race and Sex Group

	White			African American		
	Coefficient	Individual Example Value	Coefficient × Value†	Coefficient	Individual Example Value	Coefficient × Value†
Women (Example: 55 years of age with total cholesterol 213 mg/dL, HDL-C 50 mg/dL, untreated systolic BP 120 mm Hg, nonsmoker, and without diabetes)						
Ln Age (y)	-29.799	4.01	-119.41	17.114	4.01	68.58
Ln Age, Squared	4.884	16.06	78.44	N/A	N/A	N/A
Ln Total Cholesterol (mg/dL)	13.540	5.36	72.59	0.940	5.36	5.04
Ln Age×Ln Total Cholesterol	-3.114	21.48	-66.91	N/A	N/A	N/A
Ln HDL-C (mg/dL)	-13.578	3.91	-53.12	-18.920	3.91	-74.01
Ln Age×Ln HDL-C	3.149	15.68	49.37	4.475	15.68	70.15
Log Treated Systolic BP (mm Hg)	2.019	–	–	29.291	–	–
Log Age×Log Treated Systolic BP	N/A	N/A	N/A	-6.432	–	–
Log Untreated Systolic BP (mm Hg)	1.957	4.79	9.37	27.820	4.79	133.19
Log Age×Log Untreated Systolic BP	N/A	N/A	N/A	-6.087	19.19	-116.79
Current Smoker (1=Yes, 0=No)	7.574	0	0	0.691	0	0
Log Age×Current Smoker	-1.665	0	0	N/A	N/A	N/A

Goff DC Jr, et al.
2013 ACC/AHA Cardiovascular Risk Guideline

Diabetes (1=Yes, 0=No)	0.661	0	0	0.874	0	0
Individual Sum			-29.67			86.16
Mean (Coefficient× Value)	N/A	N/A	-29.18	N/A	N/A	86.61
Baseline Survival	N/A	N/A	0.9665	N/A	N/A	0.9533
Estimated 10-Y Risk for hard ASCVD	N/A	N/A	2.1%	N/A	N/A	3.0%
Men (Example: 55 years of age with total cholesterol 213 mg/dL, HDL-C 50 mg/dL, untreated systolic BP 120 mm Hg, nonsmoker, and without diabetes)						
Log Age (y)	12.344	4.01	49.47	2.469	4.01	9.89
Log Total Cholesterol (mg/dL)	11.853	5.36	63.55	0.302	5.36	1.62
Log Age×Log Total Cholesterol	-2.664	21.48	-57.24	N/A	N/A	N/A
Log HDL-C (mg/dL)	-7.990	3.91	-31.26	-0.307	3.91	-1.20
Log Age×Log HDL-C	1.769	15.68	27.73	N/A	N/A	N/A
Log Treated Systolic BP (mm Hg)	1.797	-	-	1.916	-	-
Log Untreated Systolic BP (mm Hg)	1.764	4.79	8.45	1.809	4.79	8.66
Current Smoker (1=Yes, 0=No)	7.837	0	0	0.549	0	0
Log Age×Current Smoker	-1.795	0	0	N/A	N/A	N/A
Diabetes (1=Yes, 0=No)	0.658	0	0	0.645	0	0
Individual Sum			60.69			18.97
Mean (Coefficient× Value)	N/A	N/A	61.18	N/A	N/A	19.54
Baseline Survival	N/A	N/A	0.9144	N/A	N/A	0.8954

Goff DC Jr, et al.
2013 ACC/AHA Cardiovascular Risk Guideline

Estimated 10-Y Risk for hard ASCVD	N/A	N/A	5.3%	N/A	N/A	6.1%
--	-----	-----	------	-----	-----	------

*Defined as first occurrence of nonfatal MI or CHD death, or fatal or nonfatal stroke.

†Coefficient×Value: For age, lipids, and BP, defined as the natural log of the value multiplied by the parameter estimate. When an age interaction is present with lipids or BP, the natural log of age is multiplied by the natural log of the lipid or BP, and the result is multiplied by the parameter estimate. “N/A” indicates that that specific covariate was not included in the model for that sex-race group; “–” indicates that this value was not included in the example (e.g., this example used untreated systolic BP, not treated systolic BP).

ASCVD indicates atherosclerotic cardiovascular disease; BP indicates blood pressure; CHD, congestive heart disease; HDL-C, high-density lipoprotein cholesterol; MI, myocardial infarction; and N/A, not included.

Table B. Estimating an Individual’s 10-Year Risk for Incident Hard ASCVD

The hypothetical profile provided in Table 5 (the “Individual Example Value” column) is identical for each race and sex group and is based on the overall sample mean. The profile assumes an individual 55 years of age (for which the $\text{Ln}[\text{Age}] = 4.01$), with a total cholesterol of 213 mg/dL, HDL-C of 50 mg/dL, and an untreated systolic BP of 120 mm Hg. This individual is not a current smoker and does not have diabetes. For the equations, the values for age, lipids, and systolic BP are log transformed. Interactions between age and lipids or age and systolic BP use the natural log of each variable (e.g., $\text{Ln}[\text{Age}] \times \text{Ln}[\text{Total Cholesterol}]$).

Calculation of the 10-year risk estimate for hard ASCVD can best be described as a series of steps. The natural log of age, total cholesterol, HDL-C, and systolic BP are first calculated with systolic BP being either a treated or untreated value. Any appropriate interaction terms are then calculated. These values are then multiplied by the coefficients from the equation (“Coefficient” column of Table A) for the specific race-sex group of the individual. The “Coefficient×Value” column in the table provides the results of the multiplication for the risk profile described above.

The sum of the “Coefficient×Value” column is then calculated for the individual. For the profile shown in Table A, this value is shown as “Individual Sum” for each race and sex group.

The estimated 10-year risk of a first hard ASCVD event is formally calculated as 1 minus the survival rate at 10 years (“Baseline Survival” in Table A), raised to the power of the exponent of the “Coefficient×Value” sum minus the race and sex specific overall mean “Coefficient×Value” sum; or, in equation form:

$$1 - S_{10}^{e^{(\text{Ind}X'B - \text{Mean}X'B)}}$$

Using White men as an example:

$$1 - 0.9144^{e^{(60.59 - 61.18)}}$$

equates to a 5.3% probability of a first hard ASCVD event within 10 years.

ASCVD indicates atherosclerotic cardiovascular disease; BP, blood pressure; and HDL-C, high-density lipoprotein cholesterol.

The Work Group also considered the inclusion of additional and novel risk markers in the risk equations. Based on the availability of data across cohorts at applicable examination cycles, additional risk markers

Goff DC Jr, et al.

2013 ACC/AHA Cardiovascular Risk Guideline

were evaluated for potential inclusion if they improved model performance using the framework of Hlatky et al (40). The additional risk markers that were evaluated included diastolic BP; family history of ASCVD; moderate or severe chronic kidney disease (defined as an estimated GFR of <60 mL/min per 1.73 m^2) (65); and body mass index (continuous or categorical). None of these variables significantly improved discrimination for 10-year hard ASCVD risk prediction when added to the final base models. Other risk markers (hs-CRP, ApoB, microalbuminuria, cardiorespiratory fitness, CAC score, CIMT, and ABI) could not be evaluated in creating this new model due to absence of data or lack of inclusion in the appropriate examination cycle of 1 or more of the studies. Therefore, these and the other risk markers were addressed in CQ1 as potential adjuncts to quantitative risk estimation.

Further research using state-of-the art statistical techniques (including net reclassification improvement and integrative discrimination index (66)) are needed to examine the utility of novel biomarkers when added to these new Pooled Cohort Equations in different populations and patient subgroups. Randomized clinical trials demonstrating the utility of screening with novel risk markers would represent the best evidence for their inclusion in future risk assessment algorithms. In the absence of evidence from trials, methodologically rigorous observational studies should be conducted to evaluate utility.

Goff DC Jr, et al.
2013 ACC/AHA Cardiovascular Risk Guideline

Appendix 5. Author Relationships With Industry and Other Entities (Relevant)—2013 ACC/AHA Guideline on Assessment of Cardiovascular Risk

Work Group Member	Employment	Consultant	Speaker's Bureau	Ownership/ Partnership/Principal	Personal Research	Expert Witness
David C. Goff, Jr <i>Co-Chair</i>	Colorado School of Public Health—Dean	2008-2012: None	2008-2012: None	2008-2012: None	2008-2012: • Merck	2008-2012: None
		2013: None	2013: None	2013: None	2008-2012: None	2013: None
Donald M. Lloyd-Jones <i>Co-Chair</i>	Northwestern University Feinberg School of Medicine— Senior Associate Dean; Chair and Professor of Preventive Medicine; Professor of Medicine (Cardiology)	2008-2012: None	2008-2012: None	2008-2012: None	2008-2012: None	2008-2012: None
		2013: None	2013: None	2013: None	2013: None	2013: None
Glen Bennett <i>Ex-Officio</i>	NHLBI—Coordinator	2008-2012: None	2008-2012: None	2008-2012: None	2008-2012: None	2008-2012: None
		2013: None	2013: None	2013: None	2013: None	2013: None
Sean Coady <i>Ex-Officio</i>	NHLBI—Statistician	2008-2012: None	2008-2012: None	2008-2012: None	2008-2012: None	2008-2012: None
		2013: None	2013: None	2013: None	2013: None	2013: None
Ralph B. D'Agostino	Boston University—Professor of Mathematics and Statistics; Mathematics and Statistics Department—Chair	2008-2012: None	2008-2012: None	2008-2012: None	2008-2012: None	2008-2012: None
		2013: None	2013: None	2013: None	2013: None	2013: None
Raymond Gibbons	Nuclear Cardiology Laboratory Mayo Clinic—Professor of Medicine and Co-Director	2008-2012: None	2008-2012: None	2008-2012: None	2008-2012: None	2008-2012: None
		2013: • AstraZeneca • Lantheus Medical Imaging	2013: None	2013: None	2013: None	2013: None
Philip Greenland	Northwestern University Feinberg School of Medicine— Senior Associate Dean for Clinical and Translational Research; Harry W. Dingman	2008-2012: None	2008-2012: None	2008-2012: None	2008-2012: None	2008-2012: None
		2013:	2013:	2013:	2013:	2013:

Goff DC Jr, et al.
2013 ACC/AHA Cardiovascular Risk Guideline

	Professor of Medicine	None	None	None	None	None
Daniel T. Lackland	Medical University of South Carolina—Professor of Epidemiology and Medicine	2008-2012: None	2008-2012: None	2008-2012: None	2008-2012: None	2008-2012: None
		2013: None	2013: None	2013: None	2013: None	2013: None
Daniel Levy <i>Ex-Officio</i>	NHLBI—Framingham Heart Study, Director	2008-2012: None	2008-2012: None	2008-2012: None	2008-2012: • BG Medicine	2008-2012: None
		2013: None	2013: None	2013: None	2013: None	2013: None
Christopher O'Donnell <i>Ex-Officio</i>	NHLBI—Associate Director and Senior Investigator	2008-2012: None	2008-2012: None	2008-2012: None	2008-2012: None	2008-2012: None
		2013: None	2013: None	2013: None	2013: None	2013: None
Jennifer Robinson	University of Iowa—Professor of Epidemiology and Medicine; Director, Prevention Intervention Center	2008-2012: None	2008-2012: None	2008-2012: None	2008-2012: • Aegerion • Amarin* • Amgen* • AstraZeneca* • Daiichi-Sankyo* • Esperion • Genentech/Hoffman LaRoche* • GlaxoSmithKline* • Merck* • Sanofi-aventis/Regeneron*	2008-2012: None
		2013: None	2013: None	2013: None	2013: • Amarin* • Amgen* • AstraZeneca* • Daiichi-Sankyo* • Genentech/Hoffman LaRoche* • GlaxoSmithKline* • Merck* • Sanofi-aventis/Regeneron*	2013: None

Goff DC Jr, et al.
2013 ACC/AHA Cardiovascular Risk Guideline

J. Sanford Schwartz	University of Pennsylvania—Leon Hess Professor of Internal Medicine, Health Management and Economics	2008-2012: None	2008-2012: None	2008-2012: None	2008-2012: None	2008-2012: None
		2013: None	2013: None	2013: None	2013: None	2013: None
Susan T. Shero <i>Ex-Officio</i>	NHLBI—Public Health Advisor	2008-2012: None	2008-2012: None	2008-2012: None	2008-2012: None	2008-2012: None
		2013: None	2013: None	2013: None	2013: None	2013: None
Sidney C. Smith, Jr	University of North Carolina—Professor of Medicine; Center for Cardiovascular Science and Medicine—Director	2008-2012: None	2008-2012: None	2008-2012: None	2008-2012: None	2008-2012: None
		2013: None	2013: None	2013: None	2013: None	2013: None
Paul Sorlie <i>Ex-Officio</i>	NHLBI—Chief of Division of Epidemiology and Clinical Applications	2008-2012: None	2008-2012: None	2008-2012: None	2008-2012: None	2008-2012: None
		2013: None	2013: None	2013: None	2013: None	2013: None
Neil J. Stone	Northwestern Memorial Hospital—Bonow Professor of Medicine, Feinberg School of Medicine, Northwestern University	2008-2012: None	2008-2012: None	2008-2012: None	2008-2012: None	2008-2012: None
		2013: None	2013: None	2013: None	2013: None	2013: None
Peter W.F. Wilson	Emory Clinical Cardiovascular Research Institute—Professor of Medicine	2008-2012: • Merck • XZK	2008-2012: None	2008-2012: None	2008-2012: • Merck • LipoScience	2008-2012: None
		2013: None	2013: None	2013: None	2013: None	2013: None

This table reflects the relevant healthcare-related relationships of authors with industry and other entities (RWI) provided by the panels during the document development process (2008-2012). Both compensated and uncompensated relationships are reported. These relationships were reviewed and updated in conjunction with all meetings and/or conference calls of the Work Group during the document development process. Authors with relevant relationships during the document development process recused themselves from voting on recommendations relevant to their RWI. In the spirit of full transparency, the ACC and AHA asked Work Group members to provide updates and approve the final version of this table which includes current relevant relationships (2013).

To review the NHLBI and ACC/AHA's current comprehensive policies for managing RWI, please refer to http://www.nhlbi.nih.gov/guidelines/cvd_adult/coi-rwi_policy.htm and <http://www.cardiosource.org/Science-And-Quality/Practice-Guidelines-and-Quality-Standards/Relationships-With-Industry-Policy.aspx>.

Per ACC/AHA policy:

Goff DC Jr, et al.
2013 ACC/AHA Cardiovascular Risk Guideline

A person is deemed to have a significant interest in a business if the interest represents ownership of $\geq 5\%$ of the voting stock or share of the business entity, or ownership of $\geq \$10,000$ of the fair market value of the business entity; or if funds received by the person from the business entity exceed 5% of the person's gross income for the previous year. Relationships that exist with no financial benefit are also included for the purpose of transparency. Relationships in this table are modest unless otherwise noted.

*Significant relationship.

NHLBI indicates National Heart, Lung, and Blood Institute.

Appendix 6. ACC/AHA Expert Reviewer Relationships With Industry and Other Entities

Reviewer	Employment	Representing	Consultant	Speaker's Bureau	Ownership/ Partnership/ Principal	Personal Research	Expert Witness
Ezra A. Amsterdam	University of California (Davis) Medical Center, Division of Cardiology—Professor	ACC/AHA	None	None	None	None	None
Ralph G. Brindis	University of California, San Francisco—Department of Medicine & the Phillip R. Lee Institute for Health Policy Studies—Clinical Professor of Medicine	ACC/AHA Task Force on Practice Guidelines	None	None	None	None	None
Frederick A. Masoudi	University of Colorado, Anschutz Medical Campus—Professor of Medicine (Cardiology)	ACC/AHA	None	None	None	None	None

ACC indicates American College of Cardiology and AHA American Heart Association.

Appendix 7. Abbreviations

ABI = ankle-brachial index
ACC = American College of Cardiology
AHA = American Heart Association
ApoB = apolipoprotein B
ASCVD = atherosclerotic cardiovascular disease
BP = blood pressure
CAC = coronary artery calcium
CHD = coronary heart disease
CIMT = carotid intima-media thickness
COR = class of recommendation
CQ = critical question
CV = cardiovascular
CVD = cardiovascular disease
GFR = glomerular filtration rate
HF = heart failure
hs-CRP = high-sensitivity C-reactive protein
LOE = level of evidence
NHLBI = National Heart, Lung, and Blood Institute
NHLBAC = NHLBI Advisory Council
RWI = relationships of authors with industry and other entities
Task Force = ACC/AHA Task Force on Practice Guidelines
U.S. = United States

References

1. Committee on Standards for Developing Trustworthy Clinical Practice Guidelines, Institute of Medicine. Clinical Practice Guidelines We Can Trust: The National Academies Press, 2011.
2. Gibbons GH, Harold JG, Jessup M, Robertson RM, Oetgen WJ. The Next Steps in Developing Clinical Practice Guidelines for Prevention. *Journal of the American College of Cardiology* 2013;62:1399-1400.
3. Gibbons GH, Shurin SB, Mensah GA, Lauer MS. Refocusing the Agenda on Cardiovascular Guidelines: An Announcement From the National Heart, Lung, and Blood Institute. *Journal of the American College of Cardiology* 2013;62:1396-1398.
4. Dawber TR, Kannel WB, Lyell LP. An approach to longitudinal studies in a community: the Framingham study. *Ann N Y Acad Sci* 1963;107:539-556.
5. Fried LP, Borhani NO, Enright P et al. The Cardiovascular Health Study: design and rationale. *Annals of epidemiology* 1991;1:263-76.
6. Kannel WB, Feinleib M, McNamara PM, Garrison RJ, Castelli WP. An investigation of coronary heart disease in families. The Framingham offspring study. *Am J Epidemiol* 1979;110:281-290.
7. Friedman GD, Cutter GR, Donahue RP et al. CARDIA: study design, recruitment, and some characteristics of the examined subjects. *J Clin Epidemiol* 1988;41:1105-16.
8. Investigators TA. The Atherosclerosis risk in communities (ARIC) study: design and objectives. *Am J Epidemiol* 1989;129:687-702.
9. Buckley DI, Fu R, Freeman M, Rogers K, Helfand M. C-reactive protein as a risk factor for coronary heart disease: a systematic review and meta-analyses for the U.S. Preventive Services Task Force. *Annals of internal medicine* 2009;151:483-95.
10. Empana JP, Tafflet M, Escolano S et al. Predicting CHD risk in France: a pooled analysis of the D.E.S.I.R., Three City, PRIME, and SU.VI.MAX studies. *European journal of cardiovascular prevention and rehabilitation : official journal of the European Society of Cardiology, Working Groups on Epidemiology & Prevention and Cardiac Rehabilitation and Exercise Physiology* 2011;18:175-85.
11. Fowkes FGR, Murray GD, Butcher I et al. Ankle brachial index combined with Framingham Risk Score to predict cardiovascular events and mortality: a meta-analysis. *JAMA : the journal of the American Medical Association* 2008;300:197-208.
12. Helfand M, Buckley DI, Freeman M et al. Emerging risk factors for coronary heart disease: a summary of systematic reviews conducted for the U.S. Preventive Services Task Force. *Annals of internal medicine* 2009;151:496-507.
13. Kaptoge S, Di Angelantonio E, Lowe G et al. C-reactive protein concentration and risk of coronary heart disease, stroke, and mortality: an individual participant meta-analysis. *Lancet* 2010;375:132-40.
14. Kashani M, Eliasson A, Vernalis M, Costa L, Terhaar M. Improving Assessment of Cardiovascular Disease Risk by Using Family History: An Integrative Literature Review. *The Journal of cardiovascular nursing* 2013.
15. Moyer VA. Screening for peripheral artery disease and cardiovascular disease risk assessment with the ankle-brachial index in adults: u.s. Preventive services task force recommendation statement. *Annals of internal medicine* 2013;159:342-8.
16. Peters SA, den Ruijter HM, Bots ML, Moons KG. Improvements in risk stratification for the occurrence of cardiovascular disease by imaging subclinical atherosclerosis: a systematic review. *Heart* 2012;98:177-84.
17. Schnell-Inderst P, Schwarzer R, Göhler A et al. Prognostic value, clinical effectiveness, and cost-effectiveness of high-sensitivity C-reactive protein as a marker for major cardiac events in asymptomatic individuals: a health technology assessment report. *International journal of technology assessment in health care* 2010;26:30-9.
18. Den Ruijter HM, Peters SA, Anderson TJ et al. Common carotid intima-media thickness measurements in cardiovascular risk prediction: a meta-analysis. *JAMA : the journal of the American Medical Association* 2012;308:796-803.
19. Karp I, Abrahamowicz M, Bartlett G, Pilote L. Updated risk factor values and the ability of the multivariable risk score to predict coronary heart disease. *American journal of epidemiology* 2004;160:707-16.
20. Pencina MJ, D'Agostino RB, Larson MG, Massaro JM, Vasan RS. Predicting the 30-year risk of cardiovascular disease: the framingham heart study. *Circulation* 2009;119:3078-84.
21. Lloyd-Jones DM, Leip EP, Larson MG et al. Prediction of lifetime risk for cardiovascular disease by risk factor burden at 50 years of age. *Circulation* 2006;113:791-8.

Goff DC Jr, et al.

2013 ACC/AHA Cardiovascular Risk Guideline

22. Lloyd-Jones DM, Wilson PWF, Larson MG et al. Framingham risk score and prediction of lifetime risk for coronary heart disease. *The American journal of cardiology* 2004;94:20-4.
23. Friedman GD, Cutter GR, Donahue RP et al. CARDIA: study design, recruitment, and some characteristics of the examined subjects. *J Clin Epidemiol* 1988;41:1105-1116.
24. 27th Bethesda Conference. Matching the Intensity of Risk Factor Management with the Hazard for Coronary Disease Events. September 14-15, 1995. *Journal of the American College of Cardiology* 1996;27:957-1047.
25. Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) Final Report. *Circulation* 2002;106:3143-3421.
26. Lackland DT, Elkind MS, D'Agostino R, Sr. et al. Inclusion of Stroke in Cardiovascular Risk Prediction Instruments: A Statement for Healthcare Professionals From the American Heart Association/American Stroke Association. *Stroke* 2012;43:1998-2027.
27. Shillinglaw B, Viera AJ, Edwards T, Simpson R, Sheridan SL. Use of global coronary heart disease risk assessment in practice: a cross-sectional survey of a sample of U.S. physicians. *BMC health services research* 2012;12:20.
28. Sheridan SL, Crespo E. Does the routine use of global coronary heart disease risk scores translate into clinical benefits or harms? A systematic review of the literature. *BMC health services research* 2008;8:60.
29. Sheridan SL, Viera AJ, Krantz MJ et al. The effect of giving global coronary risk information to adults: a systematic review. *Archives of internal medicine* 2010;170:230-9.
30. Redberg RF, Benjamin EJ, Bittner V et al. AHA/ACCF 2009 performance measures for primary prevention of cardiovascular disease in adults: a report of the American College of Cardiology Foundation/American Heart Association task force on performance measures (writing committee to develop performance measures for primary prevention of cardiovascular disease): developed in collaboration with the American Academy of Family Physicians; American Association of Cardiovascular and Pulmonary Rehabilitation; and Preventive Cardiovascular Nurses Association: endorsed by the American College of Preventive Medicine, American College of Sports Medicine, and Society for Women's Health Research. *Circulation* 2009;120:1296-336.
31. Aspirin for the prevention of cardiovascular disease: U.S. Preventive Services Task Force recommendation statement. *Annals of internal medicine* 2009;150:396-404.
32. Perk J, De Backer G, Gohlke H et al. European Guidelines on cardiovascular disease prevention in clinical practice (version 2012): The Fifth Joint Task Force of the European Society of Cardiology and Other Societies on Cardiovascular Disease Prevention in Clinical Practice (constituted by representatives of nine societies and by invited experts) * Developed with the special contribution of the European Association for Cardiovascular Prevention & Rehabilitation (EACPR). *Eur Heart J* 2012;33:1635-701.
33. Anderson TJ, Grégoire J, Hegele RA et al. 2012 Update of the Canadian Cardiovascular Society Guidelines for the Diagnosis and Treatment of Dyslipidemia for the Prevention of Cardiovascular Disease in the Adult. *The Canadian journal of cardiology* 2013;29:151-167.
34. Genest J, McPherson R, Frohlich J et al. 2009 Canadian Cardiovascular Society/Canadian guidelines for the diagnosis and treatment of dyslipidemia and prevention of cardiovascular disease in the adult - 2009 recommendations. *Can J Cardiol* 2009;25:567-79.
35. Morbidity & Mortality: 2012 Chartbook on Cardiovascular, Lung, and Blood Diseases. Bethesda, MD, 2012.
36. Go AS, Mozaffarian D, Roger VL et al. Heart disease and stroke statistics--2013 update: a report from the American Heart Association. *Circulation* 2013;127:e6-e245.
37. Greenland P, Knoll MD, Stamler J et al. Major risk factors as antecedents of fatal and nonfatal coronary heart disease events. *JAMA : the journal of the American Medical Association* 2003;290:891-7.
38. Magnus P, Beaglehole R. The real contribution of the major risk factors to the coronary epidemics: time to end the "only-50%" myth. *Archives of internal medicine* 2001;161:2657-60.
39. Ioannidis JP, Panagiotou OA. Comparison of effect sizes associated with biomarkers reported in highly cited individual articles and in subsequent meta-analyses. *JAMA : the journal of the American Medical Association* 2011;305:2200-10.
40. Hlatky MA, Greenland P, Arnett DK et al. Criteria for evaluation of novel markers of cardiovascular risk: a scientific statement from the American Heart Association. *Circulation* 2009;119:2408-16.
41. Di Angelantonio E, Sarwar N, Perry P et al. Major lipids, apolipoproteins, and risk of vascular disease. *JAMA : the journal of the American Medical Association* 2009;302:1993-2000.

42. Kodama S, Saito K, Tanaka S et al. Cardiorespiratory fitness as a quantitative predictor of all-cause mortality and cardiovascular events in healthy men and women: a meta-analysis. *JAMA : the journal of the American Medical Association* 2009;301:2024-35.
43. Sniderman AD, Williams K, Contois JH et al. A meta-analysis of low-density lipoprotein cholesterol, non-high-density lipoprotein cholesterol, and apolipoprotein B as markers of cardiovascular risk. *Circulation Cardiovascular quality and outcomes* 2011;4:337-45.
44. Greenland P, Alpert JS, Beller GA et al. 2010 ACCF/AHA guideline for assessment of cardiovascular risk in asymptomatic adults: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *Journal of the American College of Cardiology* 2010;56:e50-103.
45. Cavanaugh-Hussey MW, Berry JD, Lloyd-Jones DM. Who exceeds ATP-III risk thresholds? Systematic examination of the effect of varying age and risk factor levels in the ATP-III risk assessment tool. *Preventive medicine* 2008;47:619-23.
46. Marma AK, Lloyd-Jones DM. Systematic examination of the updated Framingham heart study general cardiovascular risk profile. *Circulation* 2009;120:384-90.
47. Berry JD, Dyer A, Carnethon M, Tian L, Greenland P, Lloyd-Jones DM. Association of traditional risk factors with cardiovascular death across 0 to 10, 10 to 20, and >20 years follow-up in men and women. *The American journal of cardiology* 2008;101:89-94.
48. Håheim LL, Tonstad S, Hjermann I, Leren P, Holme I. Predictiveness of body mass index for fatal coronary heart disease in men according to length of follow-up: a 21-year prospective cohort study. *Scandinavian journal of public health* 2007;35:4-10.
49. Jousilahti P, Tuomilehto J, Rastenyte D, Vartiainen E. Headache and the risk of stroke: a prospective observational cohort study among 35,056 Finnish men and women. *Archives of internal medicine* 2003;163:1058-62.
50. Liao Y, McGee DL, Cooper RS, Sutkowski MB. How generalizable are coronary risk prediction models? Comparison of Framingham and two national cohorts. *American heart journal* 1999;137:837-45.
51. Menotti A, Lanti M. Coronary risk factors predicting early and late coronary deaths. *Heart (British Cardiac Society)* 2003;89:19-24.
52. Wannamethee SG, Shaper AG, Lennon L, Morris RW. Metabolic syndrome vs Framingham Risk Score for prediction of coronary heart disease, stroke, and type 2 diabetes mellitus. *Archives of internal medicine* 2005;165:2644-50.
53. HEALTH EPOIGFC, CHILDREN RRI, ADOLESCENTS. Expert Panel on Integrated Guidelines for Cardiovascular Health and Risk Reduction in Children and Adolescents: Summary Report. *Pediatrics* 2011;128:S213-S256.
54. MD J, DH R, CM A et al. 2013 AHA/ACC/TOS guideline for the management of overweight and obesity in adults: a report of the American College of Cardiology /American Heart Association Task Force on Practice Guidelines, and The Obesity Society. *Journal of the American College of Cardiology* 2013.
55. Stone N, Robinson J, AH L et al. 2013 ACC/AHA guideline on the treatment of blood cholesterol to reduce atherosclerotic cardiovascular risk in adults: a report of the American College of Cardiology /American Heart Association Task Force on Practice Guidelines. *Journal of the American College of Cardiology* 2013.
56. Wilson PW, D'Agostino RB, Levy D, Belanger AM, Silbershatz H, Kannel WB. Prediction of coronary heart disease using risk factor categories. *Circulation* 1998;97:1837-1847.
57. D'Agostino RB, Sr., Vasan RS, Pencina MJ et al. General cardiovascular risk profile for use in primary care: The Framingham Heart Study. *Circulation* 2008;117:743-753.
58. Assmann G, Cullen P, Schulte H. Simple scoring scheme for calculating the risk of acute coronary events based on the 10-year follow-up of the prospective cardiovascular Münster (PROCAM) study. *Circulation* 2002;105:310-5.
59. Hippisley-Cox J, Coupland C, Vinogradova Y, Robson J, May M, Brindle P. Derivation and validation of QRISK, a new cardiovascular disease risk score for the United Kingdom: prospective open cohort study. *BMJ* 2007;335:136.
60. Ridker PM, Paynter NP, Rifai N, Gaziano JM, Cook NR. C-reactive protein and parental history improve global cardiovascular risk prediction: the Reynolds Risk Score for men. *Circulation* 2008;118:2243-51.
61. Ridker PM, Buring JE, Rifai N, Cook NR. Development and validation of improved algorithms for the assessment of global cardiovascular risk in women: the Reynolds Risk Score. *JAMA : the journal of the American Medical Association* 2007;297:611-9.

Goff DC Jr, et al.

2013 ACC/AHA Cardiovascular Risk Guideline

62. Conroy RM, Pyorala K, Fitzgerald AP et al. Estimation of ten-year risk of fatal cardiovascular disease in Europe: the SCORE project. *Eur Heart J* 2003;24:987-1003.
63. D'Agostino RB, Sr., Vasan RS, Pencina MJ et al. General cardiovascular risk profile for use in primary care: the Framingham Heart Study. *Circulation* 2008;117:743-753.
64. Patel MR, Dehmer GJ, Hirshfeld JW, Smith PK, Spertus JA. ACCF/SCAI/STS/AATS/AHA/ASNC 2009 Appropriateness Criteria for Coronary Revascularization A Report by the American College of Cardiology Foundation Appropriateness Criteria Task Force, Society for Cardiovascular Angiography and Interventions, Society of Thoracic Surgeons, American Association for Thoracic Surgery, American Heart Association, and the American Society of Nuclear Cardiology Endorsed by the American Society of Echocardiography, the Heart Failure Society of America, and the Society of Cardiovascular Computed Tomography. *Journal of the American College of Cardiology* 2009;53:530-553.
65. Levey AS, Stevens LA, Schmid CH et al. A new equation to estimate glomerular filtration rate. *Annals of internal medicine* 2009;150:604-12.
66. Pencina MJ, D'Agostino RB, Sr., D'Agostino RB, Jr., Vasan RS. Evaluating the added predictive ability of a new marker: from area under the ROC curve to reclassification and beyond. *Stat Med* 2008;27:157-72; discussion 207-12.